

November, 1894

TABLET TRITURATES, HYPO-
DERMIC AND COMPRESS-
ED TABLETS, PHYSICIANS'
PRIVATE FORMULAS, OFF-
ICE SUPPLIES AND VIAL
CASES.



NEW YORK
THE FRASER TABLET
TRITURATE MANUF'G CO.
17-27, Vandewater Street

Information for Our Customers.

1. The prices on our list are for thousands. Tablet Triturates are put up one thousand in a 4 ounce, French square, glass stoppered bottle. Compressed Tablets in thousands are put up in bottles ranging from 4 to 32 ozs., according to the size of tablet.

2. For five hundred Tablets, one-half the rate per thousand is charged, and five cents per 500 additional.

3. For one hundred Tablets, where the price for 100 is not given, the price is one-tenth of the price per 1000 and ten cents per 100 additional. The more expensive have the price per 100 attached.

4. The tablets are always shipped in bottles, unless otherwise specified, and in any other container are at the risk of the customer, and are not returnable.


5. Our terms are thirty days, or 1% discount for cash in 10 days, and no deviation from the list price will be made, except for marked fluctuations in the market price of the drugs.

6. When ordering, please give full address and shipping parti

NATIONAL LIBRARY OF MEDICINE Bethesda, Maryland

Gift of

William H. Helfand



FRASER'S
SOLUBLE
Tablet Triturates
(As Suggested by DR. ROBT. M. FULLER.)
—AND—
Compressed Tablets.

THESE TABLETS, INCLUDING HYPODERMICS, ETC.,
GIVE TO THE PHYSICIAN AND DRUGGIST AN EASY,
ECONOMICAL AND ACCURATE METHOD OF
DISPENSING MEDICINES IN A COMPACT
AND PALATABLE FORM.

THE directions appended to the formulas in the following list represent the individual views of the authors of the prescriptions, or have been condensed from the authorities on therapeutics. We adopt this method of replying to the many inquiries received every day from physicians as to the composition, special uses and indications of certain tablets, and in so doing assume no responsibility other than that which attaches to a desire to afford them the assistance of such information as we gather from correspondence or conversation with their fellow-practitioners in all parts of the country.

When not otherwise stated, the dose is intended for an adult, and the correction for age may be made according to the usual tables. Where the same directions are used for tablets of two or more strengths of the same medicine, it indicates the conclusions or opinions of physicians of different experience or different classes, as to the unit of dosage to be employed in specific cases, or the variation which would naturally be governed by the circumstances of each case.

When a patient cannot swallow a tablet whole, a single one may be dissolved in a proper amount of water, or a number may be dissolved in any multiple of the amount, and the directions given as for a liquid medicine, except in the cases where the tablet contains insoluble elements. For example:—One tablet in a tablespoonful of water at a dose, or eight in eight tablespoonfuls, and of this a tablespoonful as often as necessary.

Suggestions for the use of any of our tablets or for the improvement of any directions that seem inadequate, will be gratefully received. Our endeavor is to make the list as complete as we can, and as useful as possible to physicians; and while we know that the first attempt of this kind may leave much to be desired, we feel sure that it will be helpful and suggestive, and we request the aid of the medical profession in our effort to make it of even more assistance.

Please specify Fraser's Tablets

Physicians' Net Price List.

		Per Thousand.	
ABSORBENT DYSPEPTIC	{ Pepsin.....1 gr. Charcoal.....2 gr. Sodium Bicarb...2 1-2 gr. }		\$1 85
One or two after meals, followed by another in one hour, for dyspepsia and acid stomach.			
ACETANILID	1 gr.....		75
Antipyretic analgesic.			
Give one every two hours, for fever, with pain and restlessness, in children; one every twenty minutes for adults until eight are taken, if necessary.			
ACETANILID	2 grs.....		1 00
Antipyretic.			
One every half hour or every hour until three or four are taken, for fever with pain.			
ACETANILID	3 grs.....		1 20
Give one every two hours as an antipyretic until the temperature falls; also, in the same dose, to control pain in neuralgia and muscular rheumatism.			
ACETANILID	4 grs.....		1 35
One, repeated in two hours, followed by another if necessary, for neuralgia, headache and to control fever.			
ACETANILID	5 grs.....		1 50
One, repeated in two hours, followed by another if necessary, for neuralgia, headache, and to control fever.			
ACETANILID ALKALINE	{ Acetanilid.....3 gr. Spts. Ammon. Aro. 5 min. }		2 00
One every hour for three doses, to control the fever and pain of polyarthritis; then every three hours.			
ACETANILID AROMATIC	{ Acetanilid.....5 gr. Oil Gaultheria.....q. s. Pink Coloring.....q. s. }		1 65
Antipyretic.			
Give one every two hours as an antipyretic in typhoid, bronchitis and pneumonia; after the temperature falls, two to four a day.			
ACETANILID COMP. No. 1	{ Acetanilid.....2 gr. (Dr. Pitcher.) Fl. Ext. Gelsem....1 min. }		1 30
Antipyretic, analgesic.			
One every two hours for three doses, then every two hours if necessary, for neuralgia and headache, and also fever with pain.			
ACETANILID COMP. No. 2	{ Acetanilid.....4 gr. Fl. Ext. Gelsem....2 min. }		1 85
One every three hours, for fever or restlessness accompanied by pain.			
ACETANILID COMP. No. 3	{ Acetanilid.....3 gr. Diaphoretic, antipyretic. Tully's Powder....3 gr. }		2 15
One every two to four hours, for the fever of pneumonia, bronchitis and typhoid; and as an analgesic in rheumatism.			
ACETANILID COMP. No. 5	{ Acetanilid.....2 gr. (Dr. Morgan.) Quinine Sulph.....1 gr. }		2 50
One every hour for four doses, then one every two hours until relieved. For influenza and fresh colds with severe pains in the head and body.			
ACETANILID & CAFFEINE	{ Acetanilid.....3 1-2 gr. COMP.....Caffeine.....1 gr. (Dr. Hubbard.) Sodium Brom....7 1-2 gr. }		3 00
Give two tablets for the first dose, and repeat in four hours if necessary for headache and neuralgia.			

ACETANILID & CALOMEL	{ Acetanilid.....1 gr. }	1 00
COMP.	{ Sodium Bicarb....1-2 gr. }	
(Dr. J. C. Gary.)	{ Calomel.....1-10 gr. }	

Give a child from three to six years old a tablet every hour, for from three to six doses, or until bowels act. If they do not act in six hours, a saline cathartic or an enema is indicated, then give a tablet every two to four hours. For eruptive and irritative fevers, and typhoid in the first week; or in suspected cases.

In chronic cases, one every three to six hours for children or adults.

ACETANILID MODIFIED.	{ Acetanilid.....4 gr. }	1 65
	{ Tr. Digitalis.....2 min. }	

One tablet every two or three hours, for the fever and pain of pericarditis, pneumonia and bronchitis.

ACETANILID SALICYLATED.	{ Acetanilid.....4 1-2 gr. }	1 60
	{ Sodium Salicyl...1-4 gr. }	
	{ Sodium Bicarb....1-4 gr. }	

One every hour for two or three doses, then every three hours, for acute articular rheumatism and its complications

ACETANILID & SALOL	{ Acetanilid.....2 1-2 gr. }	4 00
Antiseptic, antipyretic.	{ Salol.....2 1-2 gr. }	

One every hour until fever and pain abate, in acute articular rheumatism, and neuralgia of rheumatic origin.

ACETANILID & SODA	{ Acetanilid.....4 gr. }	1 40
	{ Sodium Bicarb.....1 gr. }	

One, repeated in two hours, followed by another if necessary, for headache and neuralgia.

ACID ARSENOUS	1-1000 gr.....	75
Tonic alterative, antimalarial.		

One tablet every two hours, for an infant, as a tonic, in chronic diarrhoea, and those forms of pain and headache which have their origin in malaria.

ACID ARSENOUS	1-500 gr.....	75
----------------------	---------------	----

One to three tablets every two hours, for eczema, psoriasis and other skin diseases; also for a general tonic, in malaria and those forms of headache and neuralgia of malarial origin.

ACID ARSENOUS	1-200 gr.....	75
----------------------	---------------	----

One to two tablets every two hours, for eczema, psoriasis and other skin diseases; also for a general tonic, in malaria and those forms of headache and neuralgia of malarial origin.

ACID ARSENOUS	1-100 gr.....	75
----------------------	---------------	----

One tablet after each meal, increased by one tablet a day until effect is produced, or until fulness of the stomach with pain occurs; for a general and malarial tonic, skin diseases and those forms of neuralgic pain of malarial origin.

ACID ARSENOUS	1-60 gr.....	75
----------------------	--------------	----

One to three tablets after each meal; stop when there are colicky pains or fulness in the stomach; for skin diseases, neuralgia, epilepsy and for a malarial and general tonic.

ACID ARSENOUS	1-50 gr.....	75
----------------------	--------------	----

One to two tablets after each meal, stop when there are colicky pains or fulness in the stomach; for skin diseases, neuralgia, epilepsy and for a malarial and general tonic.

ACID ARSENOUS	1-40 gr.....	75
----------------------	--------------	----

One after each meal, increased very gradually to two after each meal, for tonic in anæmia.

ACID ARSENOUS	1-30 gr.....	75
----------------------	--------------	----

One tablet after each meal; stop when there are colicky pains or fulness in the stomach. For skin diseases, neuralgia, epilepsy and for a malarial and general tonic.

ACID ARSENOUS	1-20 gr.....	75
One three times a day after meals, for tonic in anæmia.		
ACID ARSENOUS	1-10 gr.....	75
One tablet after each meal, given with care; stop when there are colicky pains or fullness in the stomach; for skin diseases, neuralgia, epilepsy and for a malarial and general tonic.		
ACID BENZOIC	1-10 gr.....	75
Dissolve eight in half a glass of water, and give a teaspoonful every fifteen minutes for a baby, with simple or tubercular diarrhœa.		
ACID BENZOIC	1-5 gr.....	75
One every half hour in summer diarrhœa, of infants; for aphthous and diphtheritic affections of the mouth, three or four an hour, dissolved in a teaspoonful of water.		
ACID BENZOIC	1-4 gr.....	75
One every half hour for bowel complaints of children, occurring from improper food and during the summer; three to eight a day, combined with dilute hydrochloric acid, for catarrhal inflammation of the genito-urinary tract.		
ACID BENZOIC	1-2 gr.....	1 00
A tablet every hour for summer diarrhœa, cholera infantum, and entero-colitis; every half hour for diphtheria, with insufflations of powdered benzoic acid; useful also in whooping cough and bronchitis.		
ACID BENZOIC	1 gr.....	1 10
A tablet every hour for diseases of the respiratory passages, erysipelas, diphtheria, typhoid fever and acute rheumatism.		
ACID BORIC	1-10 gr.....	75
Antiseptic.		
Dissolve ten in half a glass of water, and give a teaspoonful every fifteen to thirty minutes, for an infant with diarrhœa or dysentery, and flatulence.		
ACID BORIC	1-4 gr.....	75
One every hour for infantile diarrhœa; every half hour for thrush and diphtheria, with gargle or insufflation of the powdered boric acid.		
ACID BORIC	1-2 gr.....	75
A tablet after each meal, increased to five or six a dose, or for catarrh of the ureters and bladder, and the uric acid diathesis.		
ACID BORIC	1 gr.....	95
One every hour, dissolved on the tongue; for aphthæ and diphtheria, and for diarrhœa of children.		
ACID BORIC	3 grs.....	1 15
One every two hours for dysmenorrhœa, and vesical catarrh; three or four times a day as an intestinal disinfectant.		
ACID BORIC	5 grs.....	1 30
A tablet three or four times a day, for intestinal dyspepsia with diarrhœa, and in catarrhal inflammations of the urinary tract.		
ACID CARBOLIC	1-10 gr.....	75
One every hour for indigestion with nausea, and as an intestinal disinfectant in diarrhœa, dysentery and typhoid fever.		
ACID SALICYLIC	1 gr.....	1 10
Anti-rheumatic, antipyretic.		
One every hour for three doses, then every two or three hours, for acute rheumatism of children; discontinue the medication if nausea and vomiting occur.		
ACID SALICYLIC	3 grs.....	2 00
One tablet every hour for three doses, then every three hours; for acute rheumatism and its complications, neuralgia, tonsillitis and pleurisy.		

When ordering or prescribing

ACID SALICYLIC.....5 grs..... 2 75

A tablet three or four times a day as an antipyretic; alternated with quinine to great advantage; used also in neuralgic affections, typhoid and hectic fever.

ACONITIA.....1-1000 gr..... 75

ACONITIA.....1-500 gr..... 75

ACONITIA.....1-400 gr..... 1 00

ACONITIA.....1-200 gr..... 2 00

ACONITIA.....1-150 gr..... 3 00

ACONITIA.....1-100 gr..... 4 00

ACONITIA.....1-50 gr..... 6 00

ACONITE & BELLADONNA { Aconite Powd..1-100 gr. }
 { Belladon. Powd..1-100 gr. } 75

One crushed and taken on the tongue, or mixed with water on the end of a spoon, to a baby under three, at intervals of an hour, for bronchitis, pleurisy and fevers; for older children, one every half hour for four doses, then every hour.

ACONITE & BELLADONNA { Tr. Aconite,....1-2 min. }
 No. 2..... { Tr. Belladonna..1-2 min. } 75

One every hour until effect is produced. For children one tablet in ten teaspoonfuls of water, and give a teaspoonful every twenty minutes, for fever, tonsillitis and early stages of inflammatory diseases.

ACONITE & BELLADONNA { Tr. Aconite,....1 min. }
 No. 3..... { Tr. Belladonna..1 min. } 95

One every hour for measles and other fevers of children, acute coughs and colds and acute rheumatism; after the sixth dose, give every three hours.

ACONITE & BRYONIA..... { Aconite Powd..1-100 gr. }
 Sedative, antipyretic, mem- { Bryonia Powd..1-100 gr. } 75
 brane stimulant.

One every half hour as a sedative and antipyretic in the early stages of fever, and cold with fever.

ACONITE & BRYONIA, No. 2. { Tr. Aconite,....1 min. }
 { Tr. Bryonia.....1 min. } 100

One, three or four times a day, increasing carefully according to the effect; for pneumonia and for pleurisy with effusion.

ACONITE & BRYONIA COMP. { Tr. Aconite,....1 min. }
 (Dr. Judd.) { Tr. Gelsem,....1 7-8 min. }
 { Tr. Bryonia....1 1-4 min. } 1 35

Dissolve ten tablets in four ounces of water and give a teaspoonful every hour in all febrile conditions, and especially in those due to inflammation of the air-passages.

ACONITE COMP..... { Morph. Sulph....1-50 gr. }
 { Tartar Emetic...1-100 gr. }
 { Ext. Aconite....1-100 gr. } 1 25

One every three hours; for a child with bronchitis and high fever.

ACONITE FL. EXT.....1 minim..... 75
 Diaphoretic.

Dissolve one in six tablespoonfuls of water; of this give one table- spoonful every hour, in neuralgia and fever.

ACONITE & GELSEMIUM..... { Tr. Aconite,....2 min. }
 { Tr. Gelsemium...1 min. }
 { Tr. Belladonna....2 min. } 1 50

Give one in water, for sick headache and neuralgia; repeat hourly for two doses if necessary.

ACONITE & IPECAC.	{ Tr. Aconite..... 1-2 min. }	1 00
Sedative, diaphoretic, expectorant.	{ Wine of Ipecac.... 1 min. }	

One every half hour for five doses, then one every hour or two, as a sedative and diaphoretic; for colds with hoarseness and dryness of the throat.

ACONITE & MERCURIUS.	{ Aconite Powd... 1-100 gr. }	75
Sedative, alterative.	{ Merc. Sol. H.... 1-100 gr. }	

One every two hours for a child, for fever or cold with fever and congestion.

ACONITE RAD. PULV.	1-200 gr.....	75
Antipyretic sedative.		

One every hour in the fevers of children, when pulse is rapid and of high tension.

ACONITE RAD. PULV.	1-100 gr.....	75
--------------------------------	---------------	----

Give one tablet every hour or two, for elevated temperature, strong pulse and restlessness, and for colds of children, with fever.

ACONITE RAD. PULV.	1-10 gr.....	75
--------------------------------	--------------	----

One tablet every two or three hours, for colds with fever.

ACONITE & STIBIUM.	{ Aconite Powd... 1-50 gr. }	75
(Dr. Wood.)	{ Tartar Emetic... 1-50 gr. }	
Sedative expectorant.		

One or two every half hour for four doses; repeat after two hours, in cases of dry and obstinate cough, asthma and chronic bronchitis.

AGARICIN	1-10 gr.....	1 50
-----------------------	--------------	------

One or two at night, for the profuse sweating of phthisical and gouty patients, and for exhausting cough.

AIKEN'S TONIC	{ Quinine Sulph..... 1 gr. }	2 00
	{ Iron Powd..... 2-3 gr. }	
	{ Acid Arsenous... 1-50 gr. }	
	{ Strych. Sulph..... 1-50 gr. }	
	{ Ext. Gentian..... 1-4 gr. }	

One to two tablets, three times a day, after meals.

ALKALINE CHARCOAL	{ Charcoal..... 1 gr. }	75
Absorbent alkaline.	{ Sodium Bicarb..... 1 gr. }	

Give frequently with a little water; for heartburn, acidity of the stomach and flatulence.

ALKALINE IODIDE	{ Potass. Iodide..... 5 gr. }	6 00
Alterative.	{ Potass. Bicarb..... 5 gr. }	

One after eating, for depressed and weakened conditions of the system, poverty of the blood, and chronic syphilitic conditions.

ALKALINE LAXATIVE	{ Ext. Belladonna.. 1-8 gr. }	3 00
	{ Ext. Nux Vom.... 1-4 gr. }	
	{ Ext. Colocy. Co.. 1 1-2 gr. }	
	{ Sodium Bicarb.... 1 gr. }	

One every night; for chronic constipation and the intestinal dyspepsia of rheumatic and gouty subjects.

ALOES, COLOCYNTH COMP. ..	{ Aloes..... 1 gr. }	3 25
(Dr. Duncan.)	{ Ext. Colocynth Co.. 2 gr. }	
	{ Ext. Belladonna.. 1-8 gr. }	

One tablet before dinner and two at bedtime, followed for several days by one or two at bedtime, to relieve torpidity or passive congestion of the liver

ALOIN	1-10 gr.....	75
--------------------	--------------	----

One tablet at night occasionally, for constipation of children due to errors in diet.

ALOIN	1-5 gr.....	75
--------------------	-------------	----

One or two at night in habitual constipation.

When ordering or prescribing

ALOIN	1-4 gr.....	75
One or two at night, in chronic constipation of sedentary people.		
ALOIN	1-2 gr.....	1 00
One to four tablets at night, for constipation.		
ALOIN COMP. No. 1	{ Aloin.....1-8 gr. } Cathartic hepatic. { Podophyllin.....1-8 gr. }	1 10
One to four at night, as a hepatic stimulant in chronic constipation; gradually reduce the dose.		
ALOIN COMP. No. 2	{ Aloin.....1-4 gr. } Podophyllin.....1-4 gr. }	1 25
One or two at night as a laxative, where the action of the liver is sluggish.		
ALOIN & BELLADONNA CO.	{ Aloin.....1-5 gr. } Ext. Belladonna...1-8 gr. } Tonic laxative. { Strych. Sulph....1-60 gr. }	1 10
One or two tablets at night, repeated every three hours the next day if necessary; for torpid liver and as a laxative in chronic constipation.		
ALOIN & BELLADONNA CO.	{ Aloin.....1-5 gr. } Ext. Belladonna...1-8 gr. } No. 2..... { Strych. Sulph....1-120 gr. }	1 10
One or two at night, to be repeated the next morning if necessary; as a tonic laxative.		
ALOIN, BELLADONNA AND NUX	{ Aloin.....1-5 gr. } Ext. Belladonna...1-8 gr. } (Dr. Willard.) { Ext. Nux Vomica 1-6 gr. }	1 25
One every night, decreasing to one every second night, in cases of habitual constipation.		
ALOIN, BELLADONNA AND PODOPHYLLIN	{ Aloin.....1-8 gr. } Ext. Belladonna...1-8 gr. } (Dr. LeMoyne.) { Podophyllin.....1-8 gr. }	1 25
Hepatic, tonic laxative.		
ALOIN, BELLADONNA AND PODOPHYLLIN COMP.	{ Aloin.....1-10 gr. } Ext. Belladonna...1-10 gr. } Podophyllin.....1-10 gr. } Ext. Nux Vom...1-10 gr. }	1 25
One or two at bedtime as a hepatic laxative, repeated every three hours the next day, as a tonic to liver; then one every second or third night for constipation.		
ALOIN, CASCARIN COMP	{ Aloin.....1-5 gr. } Ext. Belladonna...1-8 gr. } (Dr. Duncan.) { Cascarin.....1-4 gr. } Tonic hepatic. { Strych. Sulph....1-60 gr. }	1 50
ALOIN, CASCARIN CO. No. 2	{ Aloin.....1-4 gr. } Podophyllin.....1-4 gr. } (Dr. Duncan.) { Cascarin.....1-4 gr. } Ext. Belladonna...1-8 gr. }	1 60
ALOIN, HYOSCYAMUS CO.	{ Aloin.....1-4 gr. } Ext. Belladonna...1-4 gr. } (Dr. Wilkinson.) { Ext. Hyoscyamus 1-4 gr. }	1 40
One or two every night, for cases of constipation with intestinal discomfort and hemorrhoids.		
ALOIN, IRON & STRYCHNINE	{ Aloin.....1-10 gr. } (Dr. E. Jay Fisk.) { Iron Powd.....1 gr. } Hæmatic, tonic, cathartic. { Strych. Sulph....1-60 gr. }	1 10
One or two, after each meal, dose to be governed by the laxative effect, for the atonic dyspepsia and non-assimilation of the anæmic; also in anæmia and general debility with constipation.		

ALOIN, PODOPHYLLIN AND HYOSCYAMUS	{ Aloin.....1-4 gr.) Podophyllin.....1-4 gr.) Ext. Hyoscyam...1-10 gr.)	1 25
(Dr. Clark.)		

One or two at night, followed by one next morning if necessary, for constipation.

ALOIN, SAGRADIN COMP	{ Aloin.....1-8 gr.) Strych. Sulph....1-120 gr.) Cascarin.....1-6 gr.) Ext. Hyoscyam...1-4 gr.)	1 25
(Dr. Morris.)		

One or two at night, followed by one every three hours the next day until catharsis is produced; afterwards one at night as necessary; in chronic constipation and any cholæmic condition where a mild but good cholagogue is needed.

ALOIN & STRYCHNINE CO. ..	{ Aloin.....1-5 gr.) Ext. Belladonna...1-8 gr.) Strych. Sulph....1-60 gr.) Ipecac Powd....1-16 gr.)	1 10
Laxative, stomachic.		

One before supper and at bedtime, followed by one every two hours next day if necessary, for constipation, sick headache and as a tonic to the liver.

ALOIN STRYCHNINE CO. No 2	{ Aloin.....1-5 gr.) Ext. Belladonna...1-16 gr.) Strych. Sulph....1-60 gr.) Ipecac Powd....1-16 gr.)	1 10
----------------------------------	---	------

One before supper and at bedtime, followed by one every two hours next day if necessary, for constipation, sick headache and as a tonic to the liver.

ALTERANS	{ Acid Arsenous...1-20 gr.) Strych. Sulph....1-30 gr.) Merc. Bichlor....1-40 gr.) Iron Powd.....1 gr.)	1 00
(Dr. Hamilton.)		

A tonic in certain intractable nerve diseases, to be used under proper indications.

ALUM COMP	{ Alum.....2 1-2 gr.) Zinc Sulphate.....2 gr.) Ex. Hydras colorless 1 gr.) Morph. Sulph....1-32 gr.)	2 00
------------------------	---	------

Dissolve one in two to four tablespoonfuls of water for use as an injection in gonorrhœa and leucorrhœa.

AMMONIUM BROMIDE	5 grs.	1 10
Sedative.		

One or two at night, repeated in an hour, to quiet the nerves and induce sleep.

AMMONIUM BROMIDE	10 grs.	2 00
-------------------------------	---------	------

Give one tablet every two hours, repeated as many as three times, if necessary, for cerebral congestion and insomnia.

AMMONIUM MURIATE	1-10 gr.	75
------------------------------	----------	----

For an infant, one tablet crushed in a little water, every half hour, for bronchitis and difficult breathing; one every 15 to 30 minutes to induce expectoration in the adult.

AMMONIUM MURIATE	1-4 gr.	75
-------------------------------	---------	----

Give one every half hour for five doses, in the coughs and colds of children.

AMMONIUM MURIATE	1-2 gr.	75
-------------------------------	---------	----

A tablet allowed to dissolve on the tongue every twenty to sixty minutes as a gentle expectorant in dryness of the throat and slight colds.

AMMONIUM MURIATE	1 gr.	75
-------------------------------	-------	----

One on the tongue every hour or two as a stimulating expectorant in loss of voice from bronchial cough during secreting stages.

AMMONIUM MURIATE	3 grs.....	75
One tablet dissolved on the tongue, every two or three hours for hoarseness or during secreting stages of bronchitis.		
AMMONIUM MURIATE	5 grs.....	75
One tablet dissolved on the tongue, every two or three hours for hoarseness or during secreting stages of bronchitis.		
AMMONIUM MURIATE CO. ...	{ Ammon. Muriate 1-4 gr. Ext. Licorice 1-10 gr. Cubeb. Powd. 1-8 gr. }	75
Stimulant expectorant.		
One on the tongue every fifteen to thirty minutes, for cough with tickling and sore throat.		
AMMONIUM MURIATE CO. ...	{ Ammon. Muriate 1-4 gr. Ext. Licorice 1-10 gr. Cubeb. Powd. 1-8 gr. Ipecac Powd. 1-15 gr. }	75
No. 2		
Give one every hour, for coughs and colds, and in early stages of bronchitis.		
AMMONIUM MURIATE CO. ...	{ Ammon. Muriate 1-4 gr. Ext. Licorice 1-10 gr. Cubeb. Powd. 1-8 gr. Codeine 1-25 gr. }	1 15
WITH CODEINE		
Stimulant expectorant, and anodyne.		
Dissolve one on the tongue every fifteen to thirty minutes, for irritating cough with sore throat and tickling.		
AMMONIUM MURIATE AND ...	{ Ammon. Muriate... 1 gr. Tartar Emetic... 1-24 gr. Ext. Hyoscyam... 1-6 gr. }	1 10
HYOSCYAMUS COMP.		
Expectorant anodyne.		
One after each meal or oftener for painful cough with absent or scanty expectoration.		
AMMONIUM MURIATE CO. ...	{ Ammon. Muriate 1-4 gr. Ext. Licorice 1-10 gr. Cubeb. Powd. 1-8 gr. Morph. Sulph. 1-50 gr. }	1 10
WITH MORPHINE		
Anodyne, expectorant.		
One every half hour in cough, bronchial spasm and pleurodynia.		
AMMONIUM MURIATE AND ...	{ Ammon. Muriate 1-4 gr. Ext. Licorice 1-10 gr. Cubeb. Powd. 1-8 gr. Tartar Emetic... 1-60 gr. }	75
TARTAR EMETIC		
Diaphoretic expectorant.		
Give one, three or four times daily, in the early stages of inflammation of the respiratory passages.		
AMMONIUM SALICYLATE	5 grs.....	5 75
One tablet every three hours, for affections of the mucous membranes and secondary rheumatic manifestations.		
ANÆMIA	{ Hæmogallol,..... 3 gr. Ox Gall..... 1 gr. Ext. Pancreatis.... 3 gr. Strych. Sulph. 1-40 gr. Caffeine Mur.... 1-2 gr. Ext. Colocynth.... 1-2 gr. }	2 50
(Dr. Porter.)		
Per 100.....		
Two tablets, three times a day after meals, for one day; then one tablet after meals.		
ANÆSTHETIC	{ Camphor..... 1-4 gr. Morphine Mur.... 1-24 gr. Oil Cajeput..... 1-24 gr. }	1 50
One every hour for diarrhœa.		

ANALGESIC (Dr. Kenyon.)	{ Phenacetine.....2 1-2 gr. }	
	{ Morph. Sulph.....1-16 gr. }	
	{ Tr. Aconite.....1 min. }	
	{ Fl.Ext.Bryonia..1 1-2 min. }	
	{ Fl.Ext.Hyoscy.2 1-2 min. }	
	Per 100.....	1 50

Give two at a dose if in severe pain, otherwise one every hour for four doses, then one every two to four hours; for rheumatism, neuralgia and grip.

ANAPHRODISIAC (Dr. Nelson.)	{ Camphor.....2 gr. }	1 75
	{ Ext. Hyoscyamus...1 gr. }	

One tablet morning and night.

ANODYNE	{ Camphor.....1-8 gr. }	1 10
	{ Ext. Hyoscyam...1-8 gr. }	
	{ Morph. Sulph....1-60 gr. }	
	{ Oil Capsicum....1-60 gr. }	

One tablet every half hour until drowsiness occurs; for insomnia, restlessness and neuralgia.

ANODYNE No. 2 (Dr. H. P. Loomis.)	{ Ext. Hyoscyam.....2 gr. }	3 50
	{ Camphor.....1 gr. }	
	{ Morph. Sulph.....1-6 gr. }	

One, repeated every hour for not more than three doses, in cases of intense pain, neuralgia and tic-doloureux, intestinal and uterine colic.

ANODYNE No. 3 (Dr. Clemens.)	{ Camphor.....1 gr. }	2 25
	{ Ext. Hyoscyamus...1 gr. }	
	{ Morph. Acetate..1-20 gr. }	
	{ Oleoresin Capsic.1-20 gr. }	

One every half hour while in pain; stop after the fourth dose.

ANTACID	{ Calcium Carb....3 1-2 gr. }	1 00
	{ Magnesia Carb..2 1-2 gr. }	
	{ Sodium Chlor.....1 gr. }	

One with a little water as required, for pyrosis and stomach indigestion.

ANTIASTHMATIC	{ Potass. Iodide.....2 gr. }	4 50
	{ Potass. Bromide...2 gr. }	
	{ Fl. Ext. Euphor. Pil.3 min. }	
	{ Tinct. Lobelia...2 min. }	
	{ Nitroglycerin...1-200 gr. }	

One tablet every three or four hours, for paroxysmal cough, bronchial spasm and dyspnoea.

ANTICONSTIPATION	{ Ext. Cascara Sag...1 gr. }	2 10
	{ Ext. Nux Vom....1-8 gr. }	
	{ Ext. Belladonna..1-8 gr. }	
	{ Ipecac Powd.....1-8 gr. }	
	{ Podophyllin.....1-8 gr. }	

One to three at night, or in divided doses through the day; for sluggishness of the intestinal glands and loss of tone in the muscular coat.

ANTICONSTIPATION No. 2 (Dr. Dalton.)	{ Rhubarb Powd....1 gr. }	1 50
	{ Aloes.....1-2 gr. }	
	{ Jalap.....1-2 gr. }	
	{ Ipecac Powd.....1-4 gr. }	
	{ Castile Soap.....1-4 gr. }	
	{ Strych. Sulph....1-100 gr. }	

One or two at night.

ANTIDYSPEPTIC (Dr. Fothergill.)	{ Strych. Sulph....1-120 gr. }	1 10
	{ Ipecac Powd.....1-3 gr. }	
	{ Black Pepper Po..1-8 gr. }	
	{ Oil Gaultheria...1-10 gr. }	

One tablet after eating, repeated at the end of the hour; as a stimulant in retarded and deficient activity of the stomach.

ANTIDYSPEPTIC No. 2	{ Strych. Sulph..... 1-60 gr. }	1 25
	{ Ipecac Powd..... 1-10 gr. }	
	{ Capsicum Powd... 1-4 gr. }	
	{ Ext. Rhei..... 1-4 gr. }	
	{ Ext. Gentian..... 1-2 gr. }	
	{ Sodium Bicarb.... 2 gr. }	

One tablet in water half an hour before meals to stimulate the appetite, and repeat half an hour after eating, to hasten digestion.

ANTIDYSPEPTIC COMP. (Dr. Bradley.)	{ Cubebs Powd..... 5-8 gr. }	2 25
	{ Rhubarb Powd... 1-2 gr. }	
	{ Bismuth Subnit.... 2 gr. }	
	{ Sugar..... 2 1-2 gr. }	
	{ Oil Peppermint.. 1-4 min. }	

Give one after meals, for heartburn, eructations and deficient secretion of gastric juice.

ANTIFEBRIN COMP. (Dr. Hoag.)	{ Caffeine Citrate... 1-2 gr. }	3 75
	{ Sodium Bromide.... 5 gr. }	
	{ Antifebrin..... 2 gr. }	

One tablet with fifteen drops aromatic spts. ammonia in water, and repeat every half hour if necessary, until four doses have been taken.

ANTIFERMENTATIVE	{ Sodium Salicylate... 2 gr. }	2 00
	{ Ginger Powd..... 2 gr. }	
	{ Capsicum Powd... 1-10 gr. }	
	{ Cardamom Powd.. 1-4 gr. }	

One tablet every two or three hours for diarrhoea, flatulence and dysentery, and for the chronic intestinal dyspepsia of neurotic patients.

ANTIMALARIAL	{ Nux Vom. Powd... 1-4 gr. }	5 25
	{ Capsicum Powd... 1-2 gr. }	
	{ Ext. Hyoscyam... 1-2 gr. }	
	{ Quinine Sulph..... 3 gr. }	

Give a tablet every two or three hours in dumb-ague and general malarial disorders; give one at six, four and two hours before the chills in acute attacks.

ANTIMALARIAL No. 2	{ Cinchonidia..... 3 gr. }	
	{ Capsicum Powd... 1-2 gr. }	
	{ Nux Vomica..... 1-5 gr. }	

Give one after each meal in malaria of delicate women.

ANTIMONY & IPECAC	{ Tartar Emetic... 1-100 gr. }	75
	{ Ipecac Powd.... 1-100 gr. }	

One every two hours in the secondary stages of a cold, and as a remedy for asthma of the "dry" form.

ANTIPYRINE..... 1 gr.....	7 00
---------------------------	------

One tablet every hour for a child; stop administration as temperature falls. For adults, five every hour with a glass of water, for three doses if necessary, for headache or fever.

ANTIPYRINE..... 2 grs.....	11 50
Antipyretic. Per 100.....	1 25

Give a tablet every hour or two for headaches and fever.

ANTIPYRINE..... 3 3-4 grs.....	19 25
Per 100.....	2 00

Give one every two or three hours for neuralgic headache and headache from colds.

ANTIPYRINE..... 5 grs.....	27 00
Per 100.....	2 75

One tablet with a glass of water, every two or three hours while fever is high; for headache, one tablet every hour, for three doses if necessary.

Please specify Fraser's Tablets

ANTIPYRINE COMP.	Antipyrine.....5 gr. {	28 00
Antipyretic.	Acetanilid.....1 gr. }	
	Per 100.....	3 00

One tablet every two hours, while the fever is high; stop after the fourth dose.

ANTIRHACHITIC.	{ Precip. Chalk.....3 gr. {	1 25
(Dr. L. Weber.)	Calc. Phosph.....1-2 gr. }	
	Iron Lactate.....2-5 gr. }	

One tablet three times a day, after eating, as a constructive in rickets and imperfect or retarded bone formation.

ANTIRHACHITIC No. 2.	{ Precip. Chalk.....1-2 gr. {	1 00
(Dr. L. Weber.)	Calc. Phosph.....3-4 gr. }	
	Iron Lactate.....1-5 gr. }	

One to two tablets according to age, three times a day, after meals, as a constructive in rickets and imperfect or retarded bone formation.

ANTIRHEUMATIC.	{ Acid Salicylic.....3 gr. {	2 25
	Sodium Bicarb.....2 1-2 gr. }	
	Wine Colchicum.....1 min. }	

One tablet after each meal, and one at bedtime, for chronic rheumatism; give more frequently in acute cases.

ANTIRHEUMATIC No. 2.	{ Potass. Iodide.....2 gr. {	4 00
	Ext. Phytolacca.....1 gr. }	
	Resin Guaiac.....3 gr. }	
	Colechicin.....1-100 gr. }	
	Digitalia.....1-100 gr. }	

Give one every two hours, in a recent attack, until its severity abates; then one every four hours for not more than three doses; three times a day for chronic cases.

ANTISEPTIC No. 1.	{ Corrosive Sub.....7 3-10 gr. {	3 75
(White and Blue.)	Ammon. Mur.....7 7-10 gr. }	
	Per doz. screw cap bottles of 25 each, net	2 00

Dissolve one tablet in a pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC No. 2.	{ Corrosive Sub.....3 7-10 gr. {	2 20
(White and Blue.)	Ammon. Mur.....3 9-10 gr. }	
	Per doz. screw cap bottles of 25 each, net	2 00

Dissolve one tablet in one-half pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC No. 3.	{ Corrosive Sub.....7 3-10 gr. {	3 75
(White and Blue.)	Sodium Chlor.....7 7-10 gr. }	
	Per doz. bottles of 25 each, net	2 00

Dissolve one tablet in one pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC No. 4.	{ Corrosive Sub.....3 7-10 gr. {	2 20
(White and Blue.)	Sodium Chlor.....3 9-10 gr. }	
	Per doz. bottles of 35 each, net	2 00

Dissolve one tablet in one half pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC DYSPEPTIC.	{ Pepsin.....1 gr. {	5 50
	Bismuth Subnit.....5 gr. }	
	Salol.....1 gr. }	

A tablet a few minutes after eating, to check fermentation of food and the production of gas.

ANTISEPTIC DYSPEPTIC, AN- ODYNE.	{ Pepsin.....1 gr. {	7 50
	Bismuth Subnit.....5 gr. }	
	Salol.....1 gr. }	
	Cocaine Hydro.....1-12 gr. }	

One or two tablets after meals; for dyspepsia with gastric distress or pain.

ANTISEPTIC PASTILLES. 2 50
(Dr. Carl Seiler.)

For spray, mouth wash and nasal douche.

Dissolve one tablet in four tablespoonfuls of water, and use freely as a spray, mouth wash, gargle or nasal douche.

APERIENT. 1 75
(Dr. Drysdale.)

{ Rhubarb Powd. 1-4 gr. }	}	1 75
{ Aloes Powd. 1-4 gr. }		
{ Ipecac Powd. 5-12 gr. }		
{ Nux Vomica. 1-2 gr. }		

A tablet at night, repeated next morning if necessary, as a gentle laxative.

APHRODISIAC. 6 00

{ Ext. Cannab. Ind. 1-2 gr. }	}	6 00
{ Ext. Nux Vom. 1-2 gr. }		
{ Ext. Ergot Aq. 2 gr. }		

One tablet morning and night.

APOMORPHINE MUR. Cryst. 1-100 gr. 1 30

APOMORPHINE MUR. Cryst. 1-50 gr. 1 65

APOMORPHINE MUR. Cryst. 1-20 gr. 2 75

APOMORPHINE MUR. Cryst. 1-10 gr. 4 65

ARSENIC BROMIDE. 1-100 gr. 75

One tablet three times a day, as an alternative in sluggish skin diseases of children; stop when colic or fulness in the stomach appears.

ARSENIC BROMIDE. 1-50 gr. 75

Give one three times a day, to control sugar excretion in diabetes, and as a stimulant in chronic skin affections. Discontinue the treatment if stomach fulness or colic appears.

ARSENIC BROMIDE. 1-40 gr. 75

A tablet three or four times a day, for chronic eczema, psoriasis and malarial poisoning. Withhold the medicine if pain in the chest or abdomen occurs.

ARSENIC BROMIDE. 1-30 gr. 75

One every three or four hours; for chronic skin diseases, epilepsy and malarial poisoning. Discontinue the administration if pain in the chest or abdomen occurs.

ARSENIC BROMIDE. 1-20 gr. 75

One tablet every four hours given with care; the medicine to be withheld if pain in the stomach or abdomen should intervene; for chronic malarial infection, epilepsy and scaly skin diseases, where a tonic and stimulant are needed.

ARSENIC CHLOROPHOS. 1-50 gr. 75

One tablet every three hours as a general tonic, and for malaria in subacute and chronic forms; stop if unpleasant symptoms occur.

ARSENIC CHLOROPHOS. 1-20 gr. 75

One three times a day, stopping if pains in the chest or abdomen appear; for neuralgia, malaria and chronic skin diseases.

ARSENIC COMP. 1 50

{ Acid Arsenous. 1-30 gr. }	}	1 50
{ Piperin. 1-5 gr. }		

A tablet after meals, as a tonic and for malaria and headache of malarial origin.

ARSENICUM COMP. 75
(Dr. F. X. Spranger, Sr.)

{ Acid Arsenous. 1-100 gr. }	}	75
{ Merc. Sol. II. 1-100 gr. }		
{ Sulphur. 1-100 gr. }		

One tablet every three hours, for skin and mucous diseases, dependent on and resultant of a strumous or scrofulous diathesis.

ARSENIC IODIDE. 1-1000 gr. 75

Alterative.

One every three hours, for an infant with malaria or chronic disease of the skin, or atonic diarrhoea.

ARSENIC IODIDE	1-100 gr.....	75
One every three hours, as a stimulant tonic to the cutaneous functions, and for malarial headache. Decrease the frequency of the dose, if colic or pain in the chest appear.		
ARSENIC IODIDE	1-50 gr.....	75
Give one or two tablets after eating, as a general tonic in malaria, and as an alternative in diseases of the blood. Stop if pains in the chest or stomach develop.		
ARSENIC IODIDE	1-25 gr.....	75
One after each meal; stop when pains occur in chest or abdomen; for malaria, epilepsy, psoriasis and chronic skin diseases.		
ARSENIC IODIDE	1-10 gr.....	75
One tablet after meals given with care, as a general tonic and for malaria, epilepsy, psoriasis and chronic eczema.		
ARSENIC & IRON No. 1	Acid Arsenous...1-30 gr. } Tonic, hæmatic. } Iron Powd.....1 gr. }	75
Give a tablet three times a day, for anæmia and where a general tonic is indicated.		
ARSENIC & IRON No. 2	Acid Arsenous...1-60 gr. } Iron Powd.....1-5 gr. }	75
One or two after each meal, for malaria, anæmia and chlorosis. Stop on the occurrence of pain in the stomach or abdomen.		
ARSENIC & IRON No. 3	Acid Arsenous...1-100 gr. } Iron Powd.....1 gr. }	75
A tablet every two or three hours, for simple and pernicious anæmia, where a general tonic is required, and during convalescence after wasting disease.		
ARSENIC & IRON No. 4	Acid Arsenous...1-60 gr. } Iron Powd.....1 gr. }	75
One tablet after meals, for anæmia and chlorosis, malarial poisoning, and as a general tonic.		
ARSENIC RED SULPHURET	1-100 gr.....	75
Alternative.		
One three times a day, increasing one every second or third day; for affections of the skin accompanied with suppuration. Stop when the physiological effect is produced, or pain or fulness of the stomach occurs.		
ARSENIC RED SULPHURET	1-50 gr.....	75
One after each meal, gradually increasing to three after meals; for chronic eczema and psoriasis. Stop when the effect is produced, or pain or fulness of the stomach occurs.		
ARSENIC RED SULPHURET	1-40 gr.....	75
One three times daily, as a general alternative for diseases of the blood and skin. Stop when the effect is produced, or pain or fulness of the stomach occurs.		
ARSENIC RED SULPHURET	1-30 gr.....	75
A tablet after each meal, for boils and suppurative skin diseases. Stop when the effect is produced, or pain or fulness of the stomach occurs.		
ARSENIC RED SULPHURET	1-20 gr.....	75
One three times a day, in eczema, psoriasis and other chronic diseases of the skin. Stop when the effect is produced, or pain or fulness of the stomach occurs.		
ARSENIC & STRYCHNINE	Acid Arsenous...1-100 gr. } Tonic, alternative. } Strych. Sulph ...1-60 gr. }	75
One tablet after eating, three times a day; as a general tonic and alternative, especially in obscure malarial affections.		
ARTHRITIC	Sodium Salicylate...5 gr. } Tr. Cimicifuga....5 min. } Tr. Gelsemium....5 min. }	4 00
One every three hours for rheumatism.		

ASTRINGENT COMP.....	<div style="display: flex; align-items: center;"> <div style="flex: 1;"> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">{</div> <div> Ext. Hæmatoxylum. 1 gr. Tr. Opii Camph. . . . 6 min. Acid Sulph. Arom. . . 1 gr. Pulv. Aromat. 1 gr. </div> </div> </div> </div>	2 00
----------------------	--	------

One to three tablets according to the severity of the case, repeated in one or two hours; for diarrhœa and dysentery, and before the stage of collapse in cholera infantum.

ASTRINGENT WASH.....	<div style="display: flex; align-items: center;"> <div style="flex: 1;"> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">{</div> <div> Lead Acetate. 2 gr. Zinc Acetate. 2 gr. Hydrastin Mur. . . 1-20 gr. Morph. Acetate. . 1-32 gr. </div> </div> </div> </div>	2 00
----------------------	--	------

One in from four to eight tablespoonfuls of water for use as an injection in gonorrhœa, and also as a local application for sores, ulcers and moist skin diseases.

ATROPINE COMP..... (Dr. Gilbert.)	<div style="display: flex; align-items: center;"> <div style="flex: 1;"> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">{</div> <div> Atropine Sulph. 1-400 gr. } Tartar Emetic. . 1-100 gr. } </div> </div> </div> </div>	75
--------------------------------------	--	----

Respiratory corrective.

One every three hours in bronchitis with asthmatic complications.

ATROPINE SULPHATE.....	1-1000 gr.....	75
------------------------	----------------	----

One tablet every half hour for three doses, then every three hours, for an infant with fresh cold.

ATROPINE SULPHATE.....	1-500 gr.....	75
------------------------	---------------	----

For children, one tablet every three hours, for incontinence of urine recent colds, and to lessen the sweating of consumption and rheumatism.

ATROPINE SULPHATE.....	1-400 gr.....	75
------------------------	---------------	----

A tablet every half hour to relieve the pain of muscular rheumatism and neuralgia, and to control sweating.

ATROPINE SULPHATE.....	1-200 gr.....	75
------------------------	---------------	----

One every hour for four doses, then one every two hours, for a fresh cold.

ATROPINE SULPHATE.....	1-150 gr.....	90
------------------------	---------------	----

One every three hours for five doses, to allay vesical spasm and neuralgia, and in recent cold.

ATROPINE SULPHATE.....	1-120 gr.....	90
------------------------	---------------	----

One every three hours to relieve neuralgia and sweating of phthisical patients, and in fresh colds with abundant secretion.

ATROPINE SULPHATE.....	1-100 gr.....	1 00
------------------------	---------------	------

One every two hours for three doses, to allay vesical spasm and neuralgia, and in the early stages of colds.

ATROPINE SULPHATE.....	1-50 gr.....	1 25
------------------------	--------------	------

One tablet as an anodyne, repeated cautiously after an interval of three hours; also to relieve dyspnœa in inflammations of thoracic viscera.

BAPTISIA FL. EXT.....	1 minim.....	95
-----------------------	--------------	----

Diaphoretic, local astringent.

One tablet every three hours, as a diaphoretic, and to stimulate hepatic and intestinal functions.

BELLADONNA COMP..... (Dr. Rankin.)	<div style="display: flex; align-items: center;"> <div style="flex: 1;"> <div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">{</div> <div> Corrosive Sub. . 1-100 gr. } Tr. Belladonna. . . 1 min. } Ipecac Powd. . . . 1-10 gr. } </div> </div> </div> </div>	1 10
---------------------------------------	--	------

One every hour through the day, for ulcerated sore throat, first stages of tonsillitis and pharyngitis.

BELLADONNA EXTRACT.....	1-100 gr.....	75
-------------------------	---------------	----

A tablet every two hours for six doses, for an infant with diarrhœa and fever.

BELLADONNA EXTRACT.....	1-50 gr.....	75
-------------------------	--------------	----

One every half hour for six doses, for facial neuralgia.

BELLADONNA EXTRACT1-10 gr.....	75	
One every hour or two to control pain in neuralgia, profuse sweating in phthisis, and in recent colds.		
BELLADONNA EXTRACT1-8 gr.....	75	
A tablet every hour in recent inflammations of the respiratory tract, until three doses are taken, or dryness of the throat occurs.		
BELLADONNA EXTRACT ...1-5 gr.....	75	
Give one every two hours in recent colds, rhinitis or cold in the head, and to relieve neuralgia.		
BELLADONNA EXTRACT1-4 gr.....	75	
One tablet every three hours in pharyngitis, tonsillitis and neuralgia of the head and face.		
BELLADONNA FL. EXT1 minim.....	1 10	
Dissolve one tablet in ten tablespoonfuls of water, and give one tablespoonful of the solution every half hour, to a child with fever and headache, or recent cold.		
BETA-NAPHTOL1 gr.....	1 00	
One every half hour, with antipyretics, for tubercular and other ulceration of the bowels; when symptoms improve, give one tablet every two or three hours.		
BISMUTH AROMATIC	2 25	
(Dr. Parrish St. Louis.)		
<div>Bismuth Subnit.....2 gr. Paregoric.....3 min. Oil Cinnamon...1-20 min.</div>		
For a child one to three years of age, one tablet every hour until diarrhœa is controlled. For older children, two every hour; for all cases of inflammatory diarrhœa.		
BISMUTH COMP	1 20	
<div>Bismuth Subnit...1-2 gr. Cerium Oxalate..1-2 gr.</div>		
One or two, repeated every hour for six doses, or until effect is produced, for nausea and as sedative for sick headache.		
BISMUTH CARBONATE CO ...	1 50	
<div>Bismuth Subcarb....1 gr. Sodium Bicarb.....1 gr. Oil Peppermint...1-50 gr.</div>		
One every half hour, with a little water on the end of a spoon, for sour stomach of infants, with "spitting up" of fermented milk. Meanwhile give no milk to child, only hot water or barley water.		
BISMUTH, CATECHU CO	5 75	
<div>Salol.....2 1-2 gr. Bismuth Subnit..2 1-2 gr. Powd. Opium.....1-2 gr. Powd. Catechu....1-2 gr. Acid Hydrochl.Dil 1-2 gr. Oil Cinnamon ...1-8 min.</div>		
A tablet after meals and one at night, in chronic and subacute intestinal dyspepsia, and irritative diarrhœa.		
BISMUTH & CALOMEL CO ...		1 50
Alterative stomachic.		
<div>Bismuth Subnit.....1 gr. Calomel1-40 gr. Ipecac Powd....1-60 gr.</div>		
A tablet or two after meals, as a gentle alterative and tonic for the stomach, in dyspepsia from ice-water and overeating.		
BISMUTH & CHARCOAL	2 00	
Absorbent stomachic.		
<div>Bismuth Subnit.....2 gr. Charcoal.....5 gr.</div>		
One to three tablets after meals, for dyspepsia with fermentation and formation of gas.		
BISMUTH CHARCOAL CO	2 00	
<div>Bismuth Subcarb...2 gr. Charcoal.....2 gr. Sodium Bicarb ...3 gr. Oil Peppermint...1-10 gr.</div>		
One after eating, repeated in half an hour, for flatulent dyspepsia and heartburn; avoid starchy food.		

BISMUTH & IPECAC COMP. Alterative stomachic.	{ Bismuth Subnit.....1 gr. }	1 50
	{ Calomel.....1-10 gr. }	
	{ Ipecac Powd.....1-20 gr. }	
	{ Opium Powd.....1-40 gr. }	

One every fifteen to thirty minutes, for diarrhœa, to check the discharges easily and gradually, and at the same time to act as an intestinal antiseptic.

BISMUTH & IPECAC CO. No. 2	{ Bismuth Subnit... 1-2 gr. }	1 30
	{ Cerium Oxalate... 1-2 gr. }	
	{ Ipecac Powd.....1-20 gr. }	

A tablet as indicated in dyspepsia with nausea and headache.

BISMUTH & MORPHINE	{ Bismuth Subnit.....4 gr. }	3 65
	{ Morphine Sulph... 1-10 gr. }	

One to three tablets for vomiting and gastric pain, repeated if necessary in an hour.

BISMUTH OPIUM COMP	{ Bismuth Subnit.....4 gr. }	3 25
	{ Corros. Sub.....1-60 gr. }	
	{ Opium Powd.....1-4 gr. }	
	{ Oil Anise.....q. s. }	

One every two or three hours for colliquative diarrhœa.

BISMUTH SALICYLATE1 gr.....	1 50
--	------

Stomachic.

Give a tablet every hour or two, for intestinal fermentation; allow plenty of drinking water.

BISMUTH SALICYLATE2 grs.....	2 50
---	------

Give a tablet every hour or two, for intestinal fermentation; allow plenty of drinking water.

BISMUTH SALICYLATE5 grs.....	5 00
---	------

A tablet every two or three hours, for inflammations of the intestine, typhoid fever and dysentery; allow a free supply of drinking water.

BISMUTH SALICYLATE CO.	{ Bism. Salicyl. basic. 5 gr. }	7 50
	{ Salol.....2 1-2 gr. }	
	{ Charcoal.....2 1-2 gr. }	

One after each meal, for intestinal dyspepsia and flatulence, and the various neuralgias dependent thereon.

BISMUTH SALICYLATE AROMATIC	{ Bism. Salicyl. basic. 3 gr. }	4 00
	{ Creta Prep.....2 gr. }	
	{ Pulv. Aromat.....1 gr. }	
	{ Zinc Sulphocarb. 1 1-4 gr. }	

Give one tablet three or four times a day, in atony of the intestines, with loss of function.

BISMUTH SALICYLATE CO. No. 2	{ Bismuth Salicyl.....3 gr. }	6 75
	{ Salol.....3 gr. }	
	{ Sodium Bicarb.....2 gr. }	

One after meals for stomach or intestinal indigestion, accompanied by excessive flatulence or diarrhœa.

BISMUTH SALICIN COMP. (Dr. Hubbard.)	{ Bism. Subgall....2 1-2 gr. }	90
	{ Salicin.....5 gr. }	
	{ Strych. Sulph.. 1-100 gr. }	
	{ Per 100.....	

Two tablets four times a day, after meals and at bedtime, for atonic dyspepsia.

BISMUTH SUBCARBONATE .1 gr.	1 50
---	------

A tablet every hour until six are taken, for a child with vomiting, and the diarrhœa of teething.

BISMUTH SUBCARBONATE ..2 grs	2 00
---	------

One or two according to symptoms, for nausea, pain in the stomach and vomiting; repeated every half hour for three doses.

BISMUTH SUBCARBONATE5 grs.....	3 75	
One tablet, and repeat as necessary, for diarrhœa, dyspepsia, nausea and vomiting.		
BISMUTH SUBGALLATE2 grs.....	3 25	
(Dermatol.)		
One to three, according to age, three times a day, after meals, for flatulent or fermentative dyspepsia.		
BISMUTH SUBGALLATE3 grs.....	4 60	
(Dermatol.)		
One or two, three times a day, for fermentative or flatulent dyspepsia.		
BISMUTH SUBGALLATE4 grs.....	6 10	
(Dermatol.)		
One, three times a day, for flatulent or fermentative dyspepsia.		
BISMUTH SUBGALLATE5 grs.....	7 50	
(Dermatol.)		
One, three times a day, for flatulent or fermentative dyspepsia.		
BISMUTH SUBGALLATE CO. { Bismuth Subgall....5 gr. }	1 00	
(Dermatol.) Dr. Aulde. { Pepsin Pure.....1 gr. }		
Per 100		
One tablet before and after eating, for cases of slow or imperfect digestion of food in the stomach.		
BISMUTH SUBGALLATE CO. { Bism. Subgall....3 1-2 gr. }	8 75	
No. 2. (Dermatol.) Dr. Aulde. { Ext. Pancreatis...1 1-2 gr. }		
One tablet before meals, and two hours after, for intestinal indigestion.		
BISMUTH SUBNITRATE1 gr.....	1 35	
Stomachic.		
One tablet every two hours, for the vomiting and diarrhœa of teething children; give only hot water and sterilized milk in small quantities until symptoms abate.		
BISMUTH SUBNITRATE2 grs.....	1 85	
A tablet half an hour after eating, repeated every half hour for four doses, as a stomachic tonic and for intestinal dyspepsia.		
BISMUTH SUBNITRATE5 grs.....	3 25	
One, three times a day, in painful stomach dyspepsia; one every two hours, for diarrhœa.		
BISMUTH SUBNITRATE10 grs.....	5 85	
One as needed to allay nausea, vomiting, and gastralgia.		
BISMUTH & NUX VOMICA { Bismuth Subnit.....1 gr. }	1 50	
{ Ext. Nux Vom....1-8 gr. }		
One to three tablets, three times a day for atonic dyspepsia with lack of appetite.		
BITTER TONIC	{ Wine Ipecac.....1 min. }	2 00
	{ Tr. Capsicum.....2 min. }	
	{ Tr. Nux Vom.....5 min. }	
	{ Tr. Gentian Co....5 min. }	
One tablet just before eating, three times a day.		
BLADDER { Sodium Benzoate...5 gr. }	2 25	
(Dr. Noble.) { Sodium Biborate...5 gr. }		
One every two hours followed by a glass of water; for low grade cystitis with alkaline urine.		
BLANK TABLETS	White, Yellow and Pink....	50

When ordering or prescribing

BRONCHIAL	{	Ammon Mur.....1-3 gr.	}	1 00
		Ext. Licorice.....1 gr.		
		Bals. Tolu.....1-5 gr.		
		Oleores. Cubebs.....1-3 gr.		
		Henbane Powd.....1-4 gr.		
		Senega Powd.....1-5 gr.		
		Ipecac Powd.....1-50 gr.		

One tablet, slowly dissolved in the mouth, every hour or two, for bronchial irritation.

BRONCHIAL No. 2 (Dr. Wood.)	{	Ext. Hyoscyamus..1-5 gr.	}	1 50
		Morph. Sulph....1-100 gr.		
		Tartar Emetic...1-100 gr.		
		Ext. Aconite....1-100 gr.		
		Ext. Bellad.....1-100 gr.		
		Merc. Iod. Red..1-100 gr.		
		Potass. Bichrom.1-300 gr.		
		Bryonia.....1-200 gr.		
		Oil Anise.....1-20 gr.		

One or two as needed, every hour for inflammation of throat and bronchial tubes.

BRONCHITIS (Dr. Delafield.)	{	Ext. Belladonna.1-40 gr.	}	1 50
		Dover's Powd...1-10 gr.		
		Ipecac Powd....1-20 gr.		
		Quinine Sulph...1-4 gr.		

One every hour for four or five doses, then one every two hours for bronchial cough.

BRONCHITIS, Without Quinine (Dr. Delafield.)	{	Ext. Belladonna.1-40 gr.	}	1 00
		Dover's Powd...1-10 gr.		
		Ipecac Powd....1-20 gr.		

One every hour for four or five doses, then one every two hours, for bronchial cough.

BRONCHITIS No. 2, Acute (Dr. Davis.)	{	Tr. Aconite.....1-5 min.	}	1 00
		Tr. Bryonia....1-10 min.		
		Tr. Belladon...1-10 min.		
		Antimony Sul...1-100 gr.		
		Potass. Bichrom.1-100 gr.		

One every half hour until the bronchial secretions act normally, then every two hours as long as benefit is experienced, in bronchial cough and irritation; also useful in the chronic bronchitis of the aged; one tablet four times daily.

BRONCHITIS No. 3 (Dr. North.)	{	Tinct. Digitalis...1 min.	}	2 00
		Terpin Hydrate....1 gr.		
		Dover's Powd....1-2 gr.		
		Tinct. Belladon...1 min.		
		Tinct. Sanguinar..1 min.		
		Ammon. Mur.....1 gr.		
		Acid Benzoic....1-2 gr.		

One tablet, best allowed to dissolve slowly on the back of the tongue or dissolved in water and slowly swallowed, three or four times a day, for winter cough or bronchitis of the aged, or for abortive effects when given in first stages of grip.

BROWN AMMONIA	{	Ammon. Mur.....4 gr.	}	1 50
		Brown Mix.....1 teasp.		
		Oil Anise.....1-10 min.		

One tablet every three hours for a cough.

When ordering or prescribing

BROWN MIXTURE	(1 teaspoonful.)	{ Ext. Licorice.....1-10 gr. Camphor.....1-25 gr. Acid Benzoic....1-25 gr. Oil Anise.....1-25 gr. Opium Powd....1-25 gr. Tartar Emetic...1-60 gr.	1 00
Anodyne expectorant.			

One on the tongue every hour for bronchial cough. Stop when oppression of chest is relieved and secretion well established.

BROWN MIXTURE	($\frac{1}{2}$ teaspoonful.)	{ Ext. Licorice.....1-20 gr. Camphor.....1-50 gr. Acid Benzoic....1-50 gr. Oil Anise.....1-50 gr. Opium Powd....1-50 gr. Tartar Emetic...1-120 gr.	75
----------------------------	-------------------------------	---	----

Dissolve one in the mouth every half hour, stopping as the symptoms lessen and the cough becomes free; for bronchial coughs and colds.

BROWN MIXTURE COMP.	Throat.	{ Brown Mixture...1 teasp. Ammon. Mur.....1 gr.	1 45
---------------------------------	---------	--	------

A tablet every two hours, allowed to dissolve in the mouth, for coughs and colds with thick and difficult expectoration; lessen the frequency of the dose as the symptoms improve.

BROWN MIXT. CO. No. 2		{ Brown Mixture...1 teasp. Ammon. Mur.....5 gr.	1 35
------------------------------------	--	--	------

Take one tablet with water, or dissolve it in the mouth, every two or three hours; for bronchitis and severe cough with scanty sputa; continue until the bronchial secretions are restored.

BRUCIA	1-100 gr	Tonic.	75
---------------------	----------	--------	----

One to two tablets, three times a day, for a stimulant to the stomach and a general nerve tonic.

BRUCIA	1-50 gr		75
---------------------	---------	--	----

One tablet three times a day, for a stimulant to the stomach and a general nerve tonic.

BRUCIA COMP.	(Dr. Wood.)	{ Quinine Sulph....1-2 gr. Brucia.....1-100 gr. Acid Arsenous...1-100 gr.	1 75
Tonic antiperiodic.			

Give one after each meal as a general tonic.

BRYONIA ALB	1-200 gr	Membranous stimulant.	75
--------------------------	----------	-----------------------	----

One to three tablets, repeated every two hours as necessary, for infantile pleurisy and bronchitis.

BRYONIA ALB	1-100 gr		75
--------------------------	----------	--	----

Give one or two tablets every hour in bronchitis and pleurisy.

CACTUS COMPOUND	{ Cactus Fl. Ext....1 min. Tr. Strophanthus. 2 min.		2 35
------------------------------	--	--	------

A tablet every three or four hours, according to the severity of the symptoms, in cardiac weakness and irregularity, and as a diuretic.

CACTUS FL. EXTRACT	1 minim	Heart stimulant.	1 70
---------------------------------	---------	------------------	------

One or two tablets three times daily, as a heart stimulant in weak and irregular action from valvular disease, or myocardium.

CAFFEINE	1-10 gr		1 00
-----------------------	---------	--	------

One every hour as a stimulant to the heart, or as a diuretic.

CAFFEINE	1-2 gr		1 25
-----------------------	--------	--	------

A tablet three times a day as a cardiac stimulant and as an antiperiodic; useful also in nephritis with partial suppression.

CAFFEINE	1 gr.....	2 00
One tablet three times a day, as a stimulant to the cardiac and the renal functions; useful in the after treatment of alcohol and opium poisoning.		
CAFFEINE CITRATE	1-10 gr.....	1 00
One every hour as a stimulant to the heart, or as a diuretic.		
CAFFEINE CITRATE	1-2 gr.....	1 25
One or two tablets every three hours as a stimulant to the heart and the kidneys, and for bronchial asthma.		
CAFFEINE CITRATE	1 gr.....	2 00
A tablet three times a day to stimulate heart action, and the excretion of urine.		
CAFFEINE CITRATE	2 grs..... ¹	3 25
Headache.		
One tablet three times a day, as a diuretic in cardiac dropsy, one every two hours for three doses, for headache.		
CAFFEINE COMP	{ Caffeine Citrate....1 gr. / Nitroglycerin...1-200 gr. }	2 00
One tablet three or four times a day, for cardiac dilatation, arteriosclerosis, and as a diuretic.		
CAFFEINE COMP. No. 2	{ Caffeine Citrate....1 gr. / Nitroglycerin...1-200 gr. / Fl.Ext.Gelsem...1-2 min. }	2 25
A tablet every three or four hours as a heart tonic and stimulant, to avert the attacks of angina pectoris and lessen their severity, and to promote diuresis.		
CALCIUM SULPHIDE	1-1000 gr.....	75
A tablet every two or three hours, for the suppurative eruptions of infancy.		
CALCIUM SULPHIDE	1-100 gr.....	75
One every two hours for discharge from the ear, boils and suppurating skin affections in children; also for conjunctivitis.		
CALCIUM SULPHIDE	1-50 gr.....	75
A tablet every hour or two for boils, carbuncles, discharges from the ear and diseases of the skin.		
CALCIUM SULPHIDE	1-20 gr.....	75
One to three after meals for boils and other suppurating diseases of the cutaneous glands.		
CALCIUM SULPHIDE	1-10 gr.....	75
One every two hours for boils, carbuncles and suppurative conditions.		
CALCIUM SULPHIDE	1-8 gr.....	75
One after meals for carbuncles, furunculosis, and acne; increase gradually until the full systemic effect is reached.		
CALCIUM SULPHIDE	1-5 gr.....	75
One to two tablets three times a day for boils, carbuncles, discharges from the ear, and suppurative conditions.		
CALCIUM SULPHIDE ²	1-4 gr.....	75
One to two tablets every three hours for boils, carbuncles, discharges from the ear, and suppurative conditions.		
CALCIUM SULPHIDE	1-2 gr.....	75
One, three times a day, for boils, carbuncles and suppurative conditions.		
CALCIUM SULPHIDE	1 gr.....	1 00
A tablet three or four times daily, for suppurating affections of the skin, carbuncles, boils and otorrhœa.		

When ordering or prescribing

CALOMEL	1-1000 gr.	75
Alterative, purgative, antiseptic		
One every hour for an infant, for hepatic and intestinal tonic.		
CALOMEL	1-500 gr.	75
One every hour for an infant, for hepatic and intestinal tonic.		
CALOMEL	1-200 gr.	75
One every hour for an infant, for hepatic and intestinal tonic.		
CALOMEL	1-100 gr.	75
One every two hours to an infant for hepatic and intestinal tonic.		
CALOMEL	1-50 gr.	75
One every hour for a child until effect is produced; for hepatic and intestinal tonic, and corrective to some forms of diarrhœa.		
CALOMEL	1-40 gr.	75
One every hour for a child until effect is produced; for hepatic and intestinal tonic and corrective of diarrhœa.		
CALOMEL	1-20 gr.	75
One every hour for a child until bowels act, as corrective to liver and laxative.		
CALOMEL	1-16 gr.	75
One every hour for a child until bowels act, as corrective to liver and laxative.		
CALOMEL	1-12 gr.	75
One every half hour for adults until bowels move or twelve are taken; for hepatic alterative and laxative in biliousness or bilious headache.		
CALOMEL	1-10 gr.	75
One every hour until the bowels move, or until ten are taken.		
CALOMEL	1-8 gr.	75
One every hour until the bowels move, or until eight are taken.		
CALOMEL	1-6 gr.	75
One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia or laxative mineral water. For biliousness and torpid liver.		
CALOMEL	1-5 gr.	75
One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia or laxative mineral water. For biliousness and torpid liver.		
CALOMEL	1-4 gr.	75
One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia or laxative mineral water. For biliousness and torpid liver.		
CALOMEL	1-2 gr.	90
One every two hours until effect is produced, followed by a saline cathartic.		
CALOMEL	1 gr.	1 00
One every three hours until bowels move freely, followed by a saline cathartic.		
CALOMEL	2 gr.	1 30
One or two tablets at night as a purge, taken with a ten grain tablet of bicarbonate of sodium, followed next morning by a saline cathartic.		
CALOMEL	5 grs.	2 00
Alterative.		
One at night with a ten grain tablet of bicarbonate of soda, followed by a dose of citrate of magnesia in the morning for a purge, and alterative to liver.		

CALOMEL 10 grs. 3 00

One at night with a ten grain tablet of bicarbonate of soda, followed by a dose of citrate of magnesia in the morning for a purge, and alterative to liver.

CALOMEL, ALOIN & PODOPHYLLIN } Calomel 1-10 gr. /
 (Dr. Underwood,) } Aloin 1-10 gr. / 1 10
 Hepatic. } Podophyllin 1-10 gr. }

One to five at night, according to the requirements of the case, in chronic constipation, atony of the lower bowel, and as a hepatic stimulant; decrease the dose gradually.

CALOMEL COMP. } Calomel 1-50 gr. /
 (Dr. Spencer.) } Morph. Sulph. 1-100 gr. / 1 10
 } Tartar Emetic. 1-100 gr. }

One every hour or two until cough becomes easier.

CALOMEL COMP. No. 2 } Calomel 1-4 gr. /
 Alkaline hepatic. } Podophyllin 1-12 gr. / 75
 } Sodium Bicarb. 1-2 gr. }

One every three hours until biliousness is relieved.

CALOMEL COMP. No. 3 } Calomel 1-10 gr. /
 } Sodium Bicarb. 1 gr. / 75

One every hour until laxative effect is produced, for biliousness, headache and dull feeling with coated tongue.

CALOMEL COMP. No. 4 } Calomel 1 gr. /
 } Sodium Bicarb. 1 gr. / 1 10

One tablet at night and one in the morning, for a week; then omit the morning dose for a week, and decrease to one every second or third night. For biliousness, sick headache and light attacks of jaundice.

CALOMEL COMP. No. 5 } Calomel 1-2 gr. /
 } Podophyllin 1-12 gr. / 1 00
 } Sodium Bicarb. 1-2 gr. }

One to three at night, followed by a teaspoonful of epsom salts in cold water in the morning, as a stimulant for hepatic engorgement and to relieve hemorrhoids.

CALOMEL & BISMUTH } Calomel 1-6 gr. /
 Alterative stomachic. } Dover's Powder 1 gr. / 1 70
 } Sodium Bicarb. 2 gr. /
 } Bismuth Subnit. 1 gr. }

A tablet every three hours, as an alterative in disorders of the gastric functions.

CALOMEL & CAPSICUM } Calomel 2 gr. /
 Tonic alterative. } Capsicum 1-2 gr. / 1 50

One after meals, or two at night, follow with a glass of hot water on rising next morning. For cases of flatulent constipation and engorged portal circulation.

CALOMEL & CODEINE } Calomel 1-10 gr. /
 Sedative hepatic. } Codeine 1-8 gr. / 3 75

A tablet after meals, for cases of biliousness, with coated tongue and pain referred to the liver.

CALOMEL & IPECAC } Calomel 1-4 gr. /
 Modified hepatic and stomachic } Ipecac Powd. 1-4 gr. / 75

One or two tablets half an hour after eating, and one at bed-time, if necessary for biliousness and bilious headache, indigestion and vertigo.

CALOMEL & IPECAC No. 2 } Calomel 1-8 gr. /
 } Ipecac Powd. 1-8 gr. / 75

One every two hours for diarrhoea with green movements; for infants. Discontinue after four or five hours.

When ordering or prescribing

CALOMEL & IPECAC No. 3 ...	Calomel.....1-8 gr. }	1 10
(Dr. Nickerson.)	Ipecac Powd.....1-8 gr. }	
	Opium Powd.....1-6 gr. }	

Give one tablet every four hours, to secure the regular continuous action of mercury upon the sympathetic system.

CALOMEL & IPECAC CO.	Calomel.....1 gr. }	1 90
(Dr. Stimson.)	Ipecac Powd.....1 gr. }	
Diaphoretic alternative.	Opium Powd.....1-4 gr. }	

One at night to break up a cold; one at night until bilious stools are produced, in biliousness of patients of full habit. One every night until bowels move more freely, for feverish cold and "tight" cough.

CALOMEL & IPECAC COMP.	Calomel.....1-4 gr. }	1 35
No 2.....	Ipecac Powd.....1-2 gr. }	
	Rhubarb Powd.....1 gr. }	

One or two after each meal, with an occasional seidlitz powder in the morning; as a tonic laxative for jaundice, hemorrhoids and hepatic torpidity.

CALOMEL, IPECAC & SODA.	Calomel.....1-5 gr. }	75
Stomachic.	Ipecac Powd.....1-10 gr. }	
	Sodium Bicarb.....1 gr. }	

One or two tablets after meals, for stomach indigestion with coated tongue and headache.

CALOMEL, IPECAC & SODA	Calomel.....1-8 gr. }	75
No. 2.....	Ipecac Powd.....1-8 gr. }	
	Sodium Bicarb.....1 gr. }	

One to three tablets after eating; for dyspepsia, atony of the stomach and constipation.

CALOMEL & MORPHINE CO.	Calomel.....1-2 gr. }	3 50
Sedative hepatic.	Morphine Sulph.....1-4 gr. }	
	Tartar Emetic.....1-16 gr. }	

One tablet at night with an occasional saline laxative in the morning; for disorders of the liver accompanied by pain and neurotic symptoms.

CALOMEL & OPIUM No. 1 ...	Calomel.....1-4 gr. }	90
(Dr. McEwen.)	Opium Powd.....1-4 gr. }	

One every two hours until a laxative action is produced; for bilious colic, and as a hepatic stimulant.

CALOMEL & OPIUM No. 2	Calomel.....1-8 gr. }	75
(Dr. McEwen.)	Opium Powd.....1-8 gr. }	

One every hour until cathartic action is produced; for biliousness, headache and constipation.

CALOMEL & OPIUM CO. No. 1	Calomel.....1-2 gr. }	1 25
	Opium Powd.....1-8 gr. }	
	Ipecac Powd.....1-4 gr. }	

One every three hours, followed next morning by a saline laxative, for biliousness, jaundice and the early stages of cirrhosis.

CALOMEL & OPIUM CO. No. 2	Calomel.....1-4 gr. }	75
	Opium Powd.....1-16 gr. }	
	Ipecac Powd.....1-8 gr. }	

One every two or three hours, for bilious disorders and sick headache, followed by a saline laxative.

CALOMEL & PODOPHYLLIN.	Calomel.....1-8 gr. }	75
(Dr. Steenberg.)	Podophyllin.....1-8 gr. }	

One or two at night, or one after meals until the bowels are freely opened, for the habitual constipation of the sedentary.

CALOMEL & PODOPHYLLIN	Calomel.....1 gr. }	1 60
COMP.	Sodium Bicarb.....2 gr. }	
	Podophyllin.....1-4 gr. }	
	Ipecac Powd.....1-4 gr. }	

One three times a day, and one or two at night, for habitual constipation, plethora and tendency to piles.

CALOMEL & RHUBARB CO.	{ Calomel.....2 1-2 gr. }	2 00
Alterative, laxative.	{ Rhubarb Powd. 2 1-2 gr. }	
	{ Cinnamon Powd.1 gr. }	

One or two at night for a week, then one every second or third night, as an alterative; an occasional one at night as a laxative, for bilious tendency and intestinal dyspepsia.

CALOMEL AND SODA	{ Calomel.1-2 gr. }	95
Alkaline alterative.	{ Sodium Bicarb.3 gr. }	

One every two hours, for children's diarrhoea with green movements. Discontinue after two or three doses. Two at night for adult hepatic.

CAMPHOR COMP.	{ Camphor Monobr.1 gr. }	1 65
Analgesic.	{ Acetanilid.2 gr. }	

One every three hours when in pain; for headache, muscular rheumatism and neuralgia.

CAMPHOR, CAPSICUM CO.	{ Camphor.1-2 gr. }	2 25
Sedative stomachic.	{ Opium Powd.1-2 gr. }	
	{ Capsicum Powd.1 gr. }	
	{ Dover's Powder.2 gr. }	

A tablet half an hour after eating, for indigestion with gastralgia, headache and intestinal distress.

CAMPHOR MONOBROMIDE ..1-10 gr.....	75
Sedative.	

One every hour as a sedative in whooping cough, for a child under seven years.

CAMPHOR MONOBROMIDE ..1-2 gr.....	1 00
--	------

A tablet every hour or two, to lessen the violence of the attack in whooping cough, and quiet restlessness.

CAMPHOR MONOBROMIDE ..1 gr.....	1 35
--	------

One at night, repeated once or twice at intervals of an hour, to quiet nerves and induce sleep.

CAMPHOR MONOBROMIDE ..2 grs.....	1 80
---	------

A tablet as required to diminish the intensity of asthmatic paroxysms; also as a hypnotic and to control delirium.

CANNABIS COMP. No. 1	{ Camphor.1 gr. }	1 90
(Dr. Patterson.)	{ Ext. Can. Ind.1-4 gr. }	
	{ Ext. Hyoscyam.1-2 gr. }	
	{ Sugar of Milk.q. s. }	
Anodyne.		

Give one every two or three hours, or as often as every hour, as long as necessary to relieve pain of ovarian neuralgia, and of the genito-urinary region.

CANNABIS COMP. No. 2	{ Ext. Can. Ind.1-8 gr. }	1 70
.	{ Morph. Sulph.1-20 gr. }	
	{ Strych. Sulph.1-120 gr. }	
	{ Acid Arsenous.1-80 gr. }	
	{ Ext. Aconite.1-8 gr. }	

One tablet as required, for the pain of neuralgias or inflammations; repeat in an hour if necessary; stop at the third dose.

CANNABIS COMP. No. 3	{ Fl. Ext. Can. Ind.1 min. }	1 00
(Dr. Sadler.)	{ Fl. Ext. Gelsem.2 min. }	

One to three tablets every three to six hours, as long as pain exists. Indicated in trigeminal neuralgias and nervous disturbances arising from uterine irregularities.

CANNABIS IND. EXT1-20 gr.....	75
Hypnotic and anodyne.	

One every two hours, to control the twitchings and restlessness of chorea and induce sleep.

CANNABIS IND. EXT1-10 gr.....	1 00
--	------

A tablet after meals and one at night, for hypochondriasis and anæmic headache, and to control intense pruritus.

When ordering or prescribing

CANNABIS IND. EXT.	1-4 gr.....	1 50
Give one every three or four hours, for depression of spirits, melancholy, and some forms of headache, and to lessen the pain of rheumatism.		
CANNABIS & CODEINE	Tr. Cannabis Ind., 1 min. { Sedative. } Codeine.....1-25 gr. {	1 35
One to four tablets at intervals of half an hour, to quiet the nerves and produce sleep; in chorea, one tablet after meals.		
CANTHARIDES	1-100 gr.....	75
Stimulant diuretic.		
One tablet every hour, for congestion of urinary tract; especially to avert threatened suppression of urine; allow plenty of drinking water.		
CANTHARIDES	1-50 gr.....	75
A tablet every hour for congestion of urinary tract; administer drinking water freely.		
CANTHARIDES	1-20 gr.....	75
A tablet every three hours in renal congestion, and to prevent suppression of urine; give drinking water freely.		
CANTHARIDES	1-10 gr.....	75
Give one every three or four hours, in renal congestion with threatened anuria, with free use of drinking water.		
CAPSICUM	1-100 gr.....	75
Stomachic.		
One after meals for children, as an aid to digestion; may be repeated every hour.		
CAPSICUM	1-50 gr.....	75
One after meals for children, as an aid to digestion; may be repeated every hour.		
CAPSICUM	1-20 gr.....	75
One to three after meals, repeat one every hour afterwards, as an aid to digestion, and to relieve epigastric oppression after eating.		
CAPSICUM	1-10 gr.....	75
One or two after meals, as a tonic and aid to the digestive organs and a relief to epigastric fulness and oppression after eating. May be repeated for two doses at half hour intervals.		
CAPSICUM	1-4 gr.....	75
One or two after meals, repeated in an hour if necessary as a tonic and aid to the digestive organs, and a relief to epigastric fulness and oppression after eating.		
CAPSICUM	1-2 gr.....	75
One after each meal, repeated in an hour if necessary as a tonic and aid to the digestive organs, and a relief to epigastric fulness and oppression after eating; also good in the morning nausea of dipsomaniacs.		
CAPSICUM COMP.	{ Capsicum,.....1-10 gr. / (Dr. Lindsley.) } Nux Vomica,.....1-4 gr. }	75
Stimulant stomachic.		
One or two tablets half an hour after meals; if necessary repeat at half hour intervals for three doses, for stomach indigestion.		
CARBO VEG.	{Carbo Veg..... } (Fraser.) }Saccharin..... }	75
Antacid.		
Use freely as required, in cases of eructation, heartburn and flatulence.		
CARBO VEG.	1-100 gr.....	75
Give one every half hour to a infant with dyspepsia and wind colic.		
CARBO VEG	1-10 gr.....	75
Give one to three after meals, for dyspepsia and flatulence of children after errors in diet; give an occasional saline laxative.		

CARBO VEG	1-2 gr.	75
A tablet or two as required, for heartburn, eructation and flatulence; take a seidlitz powder in the morning every few days.		
CARDIAC	{ Sparteine Sulph. 1-10 gr. Tr. Strophanthus. 3 min. Caffeine Cit. 1-2 gr. Codeine. 1-20 gr.	3 40
(Dr. Waldstein.)		
Give a tablet every three hours, to control rapid and tumultuous heart action, irregular pulse, difficult breathing and tendency to œdema; avoid overloading the stomach.		
CARDIANE	{ Digitalia. 1-120 gr. (Merck's) Tr. Strophanthus. 2 min. Tr. Cactus. 1 min. Sparteine Sulph. 1-20 gr.	2 75
(Dr. Pallen.)		
A tablet three times a day, to regulate the action of the heart, diminish undue frequency and relieve dyspnœa; for valvular cardiac disease and begining dilatation.		
CASCARA COMP	{ Ext. Cascara. 1-2 gr. Aloin. 1-8 gr. Podophyllin. 1-10 gr. Oil Ment. Pip. 1-10 gr.	1 60
One or two at night, followed by one after each meal if necessary, for a laxative.		
CASCARA COMP. No. 2	{ Ext. Cascara Sag. 2 gr. Podophyllin. 1-8 gr. Ext. Belladonna. 1-16 gr.	2 50
Two tablets at night, as a mild tonic cathartic.		
CASCARIN COMP	{ Aloin. 1-4 gr. Podophyllin 1-4 gr. Cascarin. 1-4 gr.	1 40
Tonic, hepatic and laxative.		
A tablet after meals, with free use of drinking water, for habitual and atonic constipation.		
CASCARIN COMP No. 2	{ Ext. Cascara Sag. 1-2 gr. Aloin. 1-8 gr. Podophyllin 1-10 gr. Oil Peppermint 1-10 gr. Oil Cajeput. 1-2 gr.	1 85
(Dr. Gilbert.)		
One after each meal, until they act freely; then one at night as necessary for habitual constipation accompanied with flatulence.		
CASCARIN TABLETS	{ Salicin. 3 gr. Cascarin. 2 gr.	4 25
(Dr. Waldstein.)		
Tonic cathartic.		
One at night, for loss of tone in the intestine with consequent habitual constipation.		
CASCARA SAGRADA Fl. Ext.	5 min.	2 00
Laxative.		
One or two at night, or every three hours during the day, increasing intervals as constipation is relieved.		
CATHARTIC No. 1	{ Leptandrin. 1-32 gr. Aloin. 1-16 gr. Podophyllin 1-6 gr. Gamboge. 1-64 gr. Ext. Hyoscyam. 1-16 gr. Ol. Res. Caps. 1-128 gr. Oil Peppermint. 1-128 gr.	1 00
One to three tablets at night.		

When ordering or prescribing

CATHARTIC No. 2.....	{	Aloin.....	1-10 gr.	1 25
		Ext. Colocy. Co.....	1-10 gr.	
		Ext. Nux Vom.....	1-10 gr.	
		Podophyllin.....	1-10 gr.	
		Croton Oil.....	1-30 gr.	
		Ol. Res. Caps.....	1-256 gr.	
		Oil Peppermint.....	1-128 gr.	

One tablet every two hours until bowels move, or one to three at night, followed in the morning by another if necessary.

CATHARTIC CO. (Pill U. S. P.)..... 1-10 gr..... 75

Give one every half hour till the bowels move, or six at night, according to the requirements of the case.

CATHARTIC CO. (Pill U. S. P.)..... 1-5 gr..... 75

From one to four tablets, in divided doses through the day or at bed time; as a tonic and hepatic laxative.

CATHARTIC COMP. (Veg.)..... 1-10 gr..... 75

One every hour up to four or five doses as necessary, to procure free action of the bowels; then one every night for a time.

CATHARTIC COMP. (Veg.)..... 1-5 gr..... 75

Two or three during the day and two at night, with another in the morning if necessary; after the bowels act freely, diminish to one every night.

CATHARTIC COMP. (Imp.)..... 1-10 gr..... 75

Two tablets at night and three or four the next day; then two at night for a week, and finally every second or third night.

CATHARTIC COMP. (Imp.)..... 1-5 gr..... 75

Give one after each meal, for the gradual restoration of the activity of the bowels; then lessen the dose to one at bed time.

CAULOPHYLLIN..... 1-10 gr..... 1 10

Antispasmodic and diuretic.

Give from one to three every hour, to relieve spasm of the sphincter vesicæ and post-partum pains, and as a diuretic.

CERIUM OXALATE..... 1-10 gr..... 75

Stomachic sedative.

One every fifteen minutes for nausea of infants, diffused in water on the end of a spoon. When nausea stops, give one half an hour before each feeding for two or three meals.

CERIUM OXALATE..... 1 gr..... 1 00

One every twenty minutes for adult, every half hour for grown child; for nausea. When stomach is settled, one every two hours with hot liquid food as long as necessary.

CERIUM OXALATE..... 2 grs..... 1 40

One every hour for four or five doses, for sick stomach and nausea.

CERIUM OXALATE..... 5 grs..... 2 25

One to two tablets three or four times a day, for vomiting in pregnancy.

CERIUM OXALATE COMP...... { Cerium Oxalate..... 1 gr. /
(Dr. Guernsey.) { Ipecac Powd..... 1-20 gr. } 1 15

One every hour until nausea stops, followed by one every hour with hot beef tea in small quantities.

CHARCOAL..... 2 grs..... 75

Absorbent.

A tablet as needed to control nausea and sea-sickness, and to correct flatulence.

CHARCOAL..... 5 grs..... 75

Give one every three hours or oftener, in conditions of gastric dyspepsia with eructations.

CHARCOAL	10 grs.....	75
A tablet after each meal for indigestion, with the production of gas, nausea, and palpitation.		
CHALK	{ Chalk Prepared1 gr. { Antacid. { Oil Cinnamon... 1-20 gr. }	75
A tablet every half hour until relieved, for a child with dyspepsia and gastric distress; for a baby, give one with each feeding.		
CHALK MIXTURE	1 teasp.	75
One tablet every two or three hours, for the summer diarrhoea of children; it is best to precede this treatment with a single dose of castor oil.		
CHOCOLATE LAXATIVE	{ Ext. Cascara Sag... 1-4 gr. { Aloin.....1-40 gr. { Podophyllin.....1-40 gr. { Ext. Tamarinds.....2 gr. { Sweet Chocolate....9 gr. }	2 00
Give one to four according to age of patient, as an agreeable laxative for children.		
CHLORALAMID, Schering's	3 grs.....Per 100	1 00
A tablet three or four times a day for a child with chorea, to lessen excitability and control excessive motion; also as a hypnotic for children.		
CHLORALAMID, Schering's	5 grs.....Per 100	1 30
A tablet every three hours in chorea and to avert epileptic seizures, and as a hypnotic to replace chloral and sulphonal		
CHLORALAMID, Schering's	10 grs.....Per 100	2 50
One tablet repeated in an hour to produce sleep; for insomnia of senility, neurasthenia and alcoholism.		
CHLORALAMID, Schering's	15 grs.....Per 100	3 75
A tablet as necessary to avert epileptic and choreic attacks, to produce sleep in old age, hysteria, cardiac disease and morphinism.		
CHLORODYNE	{ Morphine Mur.... 1-6 gr. { Ext. Cannabis Ind. 1-4 gr. { Nitroglycerin... 1-300 gr. { Ext. Hyoscyam.... 1-2 gr. { Oleoresin Capsic. 1-10 gr. { Oil Ment. Pip... 1-10 gr. }	3 75
One, and repeat in an hour if necessary for pain. After taking three doses, stop three hours before resuming. For diarrhoea, one every three hours until relieved.		
CHLOROSIS	{ Quinine Sulph.....1 gr. { Iron Powd.....1 1-2 gr. { (Dr. Bishop.) { Ext. Ergot.....1 gr. { Ext. Nux Vom.... 1-5 gr. }	4 25
One tablet, three times a day, in anæmia, chlorosis and amenorrhœa.		
CHLOROSIS No. 2	{ Iron Protochlor... 1-4 gr. { Merc. Bichlor... 1-120 gr. { Arsenic Chlor.... 1-200 gr. { Quinine Chlor.... 1-8 gr. }	1 40
One or two tablets after each meal where a vigorous iron tonic is required, and during convalescence.		
CHOLAGOGUE	{ Calomel.....1 gr. { (Dr. Porter.) { Ipecac Powd.....1-8 gr. { Acid Arsenous... 1-15 gr. { Corros. Sublim... 1-15 gr. }	1 25
One every three hours until bowels move more freely; then one every second or third night, or every week, as necessary.		

When ordering or prescribing

CHOLAGOGUE No. 2	{ Calomel.....1-10 gr. }	1 00
(Dr. Kennedy.)	{ Ipecac Powd.....1-10 gr. }	
	{ Sodium Bicarb.....2 gr. }	

One every two hours, as a mild cholagogue for adults. For constipation of infants, one with as much sugar of milk as will lie on a silver ten-cent piece.

CHOLERA INFANTUM	{ Zinc Sulphocarb..1-20 gr. }	3 00
(Dr. Hillkowitz.)	{ Salol.....1-10 gr. }	
	{ Bismuth Subnit....1-2 gr. }	
	{ Calomel.....1-60 gr. }	
	{ Lactopeptin.....1-2 gr. }	

One every ten minutes until five are given, then one every hour; as a remedy for cholera infantum and summer diarrhœa in children from six to twelve months old.

CHOLERA MORBUS, Bilious..	{ Bismuth Subnit....3 gr. }	3 50
(Dr. Hitchcock.)	{ Calomel.....1-8 gr. }	
	{ Salol.....1 gr. }	

If there is an irritating substance in the bowels, give a seidlitz powder first then a tablet after each bad passage; not oftener than every hour, nor more than ten in twenty-four hours.

CIMICIFUGA FL. EXT.1 min.....	1 15
Anti-rheumatic and cardiac tonic.	

Give one to three tablets every two hours, for headache dependent upon pelvic derangement, sex hypochondriasis, and as a cardiac tonic in fatty heart and rheumatism.

COBALT & POTASS. NITRATE. 1-4 gr.....	1 10
Heart stimulant and tonic.	

A tablet every three hours as a stimulant and tonic to the cardiac function.

COBALT & POTASS. NITRATE. 1-2 gr.....	1 50
--	------

Take one after each meal, for slow and feeble action of the heart.

COCAINE COMP	{ Potass. Chlorate....1 gr. }	1 25
(Dr. F. A. Burrall.)	{ Cocaine Mur....1-100 gr. }	

One on the tongue every half hour for pharyngitis and colds.

COCAINE HYDROCHLOR1-100 gr.....	1 00
COCAINE HYDROCHLOR1-50 gr.....	1 35
COCAINE HYDROCHLOR1-20 gr.....	1 00
COCAINE HYDROCHLOR1-10 gr.....	3 00
COCAINE HYDROCHLOR1-8 gr.....	4 25
COCAINE HYDROCHLOR1-6 gr.....	5 50
COCAINE HYDROCHLOR1-5 gr.....	6 50
COCAINE HYDROCHLOR1-4 gr.....	7 75
COCAINE HYDROCHLOR1-1-8 grs.....Per 100	3 00

A tablet dissolved in a drachm of distilled water makes a two per cent. solution.

COCAINE HYDROCHLOR2 1-4 grs.....Per 100	5 50
--	------

A tablet dissolved in one drachm of distilled water makes a four per cent. solution.

COCAINE THROAT	{ Cocaine Mur.....1-12 gr. }	3 75
	{ Cubebs Powd.....1 gr. }	
	{ Acid Benzoic.....1-2 gr. }	
	{ Potass. Chlorate....2 gr. }	
	{ Licorice Powd.....q. s. }	

One on the tongue every two or three hours for chronic cough with tickling and irritation of the throat.

CODEINE	1-10 gr.....	2 75
CODEINE	1-8 gr.....	3 25
CODEINE	1-5 gr.....	4 50
CODEINE	1-4 gr.....	5 25
CODEINE	1-2 gr.....	10 50
CODEINE	1 gr.....	20 00
CODEINE PHOSPHATE	1-4 gr.....	5 75
CODEINE PHOSPHATE	1-2 gr.....	10 50
CODEINE SULPHATE	1-5 gr.....	3 75
CODEINE SULPHATE	1-4 gr.....	4 50
CODEINE SULPHATE	1-2 gr.....	8 25
COLCHICIN	1-100 gr.....	1 20

Anti-rheumatic.

One every three hours until a cathartic effect is produced. For rheumatism, gout and nephritis.

COLCHICIN	1-50 gr.....	1 70
------------------------	--------------	------

A tablet three times a day, for neuralgia, rheumatism, and chronic inflammation of the kidneys.

COLD No. 1	{ Morph. Sulph.....1-32 gr. } { Tartar Emetic.....1-32 gr. } { Ext. Licorice.....1-10 gr. } { Oil Gaultheria.....q.s. }	1 35
-------------------------	--	------

One every two or three hours, for cough with bronchial irritation; lengthen the intervals to one every four hours if nausea occurs.

COLD No. 2	{ Antim. Sul. Aur.....1-12 gr. } { Ext. Conium.....1-12 gr. } { Ipecac Powd.....1-6 gr. } { Potass. Nitrate.....1-6 gr. } { Ammon. Mur.....1-6 gr. }	1 00
-------------------------	--	------

(Dr. Satterlee.)

Two after each meal for cold of adults.

COLD No. 3	{ Quinine Sulphate.....2 gr. } { Dover's Powder.....2 gr. } { Capsicum Powd.....1-4 gr. } { Tr. Aconite.....1 min. }	4 00
-------------------------	---	------

(Dr. Gage.)

One tablet every three or four hours until effect is produced; give two or three at bedtime if necessary.

COLD No. 4	{ Aconite.....1-10 gr. } { Camphor.....1-10 gr. } { Opium Powd.....1-10 gr. } { Potass. Nitrate.....1-10 gr. }	75
-------------------------	---	----

(Dr. Coit.)

Give adults ten tablets in water at bedtime, or one every hour during the day for coryza. This treatment may be supplemented with tonic doses of quinine.

COLD No. 5. Early Stages	{ Tr. Aconite Rad.....1 min. } { Atropine Sulph.....1-600 gr. } { Morph. Sulph.....1-30 gr. } { Hydrarg. Chl. Mit. 1-6 gr. }	1 40
---------------------------------------	---	------

One every hour for three doses, then every two hours for three more; take for one day after secretion and weeping from nostrils is stopped.

COLIC	{ Morph. Sulph.....1-80 gr. } { Bismuth Subnit.....1 gr. } { Calomel.....1-40 gr. }	1 60
--------------------	---	------

(Dr. Satterlee.)

One every fifteen minutes until four are taken; rest an hour and repeat if necessary in bad cases; give an adult four at the first dose, and four every hour afterwards, as long as necessary.

When ordering or prescribing

COLIC No. 2	{ Paregoric.....2 min.	1 10
(Infantile.)	{ Sodium Bicarb.....1 gr.	
	{ Oil Fennel.....1-10 gr.	
Dissolve eight, in eight teaspoonfuls of hot water; give ten to twenty drops every ten or fifteen minutes.		
CONIUM COMP	{ Ext. Conium.....1-30 gr.	75
Expectorant.	{ Cubebs Powd....1-10 gr.	
	{ Ext. Licorice.....1-10 gr.	
One every half hour or every hour, for cough with tickling in the throat.		
CONIUM COMP. with Codeine.	{ Ext. Conium.....1-30 gr.	1 15
Anodyne expectorant.	{ Cubebs Powd....1-10 gr.	
	{ Ext. Licorice.....1-10 gr.	
	{ Codeine.....1-25 gr.	
One every hour for tickling in the throat, with persistent cough.		
CONSTIPATION. CHRONIC	{ Ext. Colocy. Co.....2 gr.	4 00
(Dr. Feltwell.)	{ Ext. Nux Vomica..1-4 gr.	
	{ Ext. Belladonna ..1-4 gr.	
	{ Ext. Gentian.....1 gr.	
	{ Oil Caraway.....1-2 gr.	
CONVALLARIA MAJAL. Fl.Ex 1 min.		1 25
Cardiac tonic.		
Two or three tablets after meals to diminish the frequency and increase the power of the heart, to remove dropsy, and to replace digitalis where that is unsafe.		
CONVALLAMARIN	1-100 gr.	1 70
One tablet three times a day, increased with great caution; for cardiac weakness and rapidity, and dropsy.		
COPAIBA BALSAM	1-100 gr.	75
Diuretic.		
One to three tablets every two hours, increasing slowly according to the severity of the case; for vesical and urethral inflammation, and to increase the flow of urine.		
COPPER ARSENITE	1-5000 gr.	75
Put five tablets in a glass of water, and give a teaspoonful every half hour, for diarrhœa in nursing infants.		
COPPER ARSENITE	1-1000 gr.	75
A tablet in a little water every two hours, for the summer diarrhœa of children.		
COPPER ARSENITE	1-500 gr.	75
Take one every hour or two until the discharges become natural.		
COPPER ARSENITE	1-200 gr.	75
A tablet three times a day for diarrhœa in children.		
COPPER ARSENITE	1-100 gr.	75
One tablet in one-half glass of water. One teaspoonful of this every half-hour until relieved of the diarrhœa.		
COPPER ARSENITE	1-50 gr.	75
One in one-half glass of water. One teaspoonful of this every half-hour until relieved of the diarrhœa.		
COPPER SULPHATE	1-120 gr.	75
One to four tablets, carefully increased, three times daily. Use with iron tonics in syphilis.		
COPPER SULPHATE	1-100 gr.	75
One after meals, with tincture ferri chlor. for syphilitic inflammations of the throat; increase after tolerance has been established.		
COPPER SULPHATE	1-60 gr.	75
One, three times a day, with iron, in the treatment of secondary syphilis.		

COPPER SULPHATE	1-40 gr.....	75
One morning and evening for secondary syphilis, best combined with iron; omit the treatment one or two days each week.		
COPPER SULPHATE	1-30 gr.....	75
One or two a day with iron, in secondary syphilis; omit treatment one or two days each week; stop for a week if the appetite becomes excessive.		
CORRECTIVE, INFANT'S	<div><div>Calomel1-20 gr.</div><div>Ipecac Powd.....1-50 gr.</div><div>Sodium Bicarb....1-2 gr.</div><div>Bismuth Subnit.....1 gr.</div></div>	1 50
(Dr. Ives.)		
A tablet, crushed and mixed with an equal bulk of sugar and placed on the child's tongue every one or two hours, to correct morbid conditions of the alimentary tract.		
CORROSIVE SUBLIMATE	1-10,000 gr.....	75
CORROSIVE SUBLIMATE	1-5000 gr.....	75
CORROSIVE SUBLIMATE	1-1000 gr.....	75
CORROSIVE SUBLIMATE	1-500 gr.....	75
CORROSIVE SUBLIMATE	1-200 gr.....	75
CORROSIVE SUBLIMATE	1-100 gr.....	75
CORROSIVE SUBLIMATE	1-60 gr.....	75
CORROSIVE SUBLIMATE	1-50 gr.....	75
CORROSIVE SUBLIMATE	1-40 gr.....	75
CORROSIVE SUBLIMATE	1-32 gr.....	75
CORROSIVE SUBLIMATE	1-30 gr.....	75
CORROSIVE SUBLIMATE	1-20 gr.....	75
CORROSIVE SUBLIMATE	1-16 gr.....	75
CORROSIVE SUBLIMATE	1-12 gr.....	75
CORROSIVE SUBLIMATE	1-10 gr.....	75
CORROSIVE SUBLIMATE	1-8 gr.....	75
CORROSIVE SUBLIMATE	1-4 gr.....	75
CORROSIVE SUBLIMATE	1-2 gr.....	90
CORROSIVE SUBLIMATE	1 gr.....	1 00
A tablet dissolved in four tablespoonfuls of water makes a solution of about 1 to 1,000; for external use.		
CORROSIVE SUB. (colored blue)	1 gr.....	1 00
Antiseptic.		
A tablet dissolved in four tablespoonfuls of water makes a solution of about 1 to 1,000; for external use.		
CORROSIVE SUBLIMATE ...	{ Corrosive Sub.1 gr. }	1 00
SOLUBLE. (For Hair Wash.)	{ Tartaric Acid.1 gr. }	
Dissolve a tablet in eight tablespoonfuls of water, and apply twice a week for dandruff.		
CORROSIVE SUBLIM. COMP. ..	{ Corrosive Sub....1-32 gr. }	75
(Dr. Cushman.)	{ Ipecac Powd.....1-8 gr. }	
Alterative.		
One tablet after meals as an alimentary stimulant and alterative, in chronic dyspepsia, rheumatism and skin diseases.		
CORROSIVE SUBLIMATE &	{ Corrosive Sub....1 3-4 gr. }	1 25
CITRIC ACID	{ Citric Acid....87-100 gr. }	
Antiseptic.		
Dissolve a tablet in four ounces of water; the strength of corrosive sublimate in the resulting solution is 1 to 1,000. The citric acid retains in solution the mercury albuminate which is formed, and thus prolongs its antiseptic action.		

CORROSIVE SUBLIMATE &	Corrosive Sub. 3-4 gr.	1 25
CITRIC ACID	Citric Acid. 87-100 gr.	
(Colored Blue.)		

Dissolve a tablet in four ounces of water, the strength of corrosive sublimate in the resulting solution is 1 to 1,000. The citric acid retains in solution the mercury albuminate which is formed, and thus prolongs its antiseptic action.

CORROSIVE SUBLIMATE &	Acid Tartaric. 5 gr.	1 50
TARTARIC ACID	Corros. Sub. 1 gr.	

One tablet in two ounces of water equals about 1 to 1,000 corrosive sublimate solution; the tartaric acid prevents the coagulation of the albumen and thus prolongs antiseptis.

CORYZA (Dr. Richards.)	Quinine Sulph.....1-2 gr.	} 2 00
	Ammon. Mur.....1-2 gr.	
	Camphor.....1-2 gr.	
	Opium Powd.....1-10 gr.	
	Ext. Belladonna.1-10 gr.	
	Ext. Aconite.....1-10 gr.	

For a fresh cold, give two tablets for first dose; followed by one every hour until cold is broken up.

CORYZA No. 2	(Quinine Sulph. 1-3 gr.)	1 50
	(Ammon. Mur. 1-3 gr.)	
	(Camphor. 1-3 gr.)	
	(Opium Powder. 1-15 gr.)	
	(Ext. Belladonna. 1-15 gr.)	
		(Ext. Aconite. 1-15 gr.)

A tablet every three hours in recent cold to prevent the spread of the inflammation and to lessen the discharge.

CORYZA No. 3	(Camphor. 1-2 gr.)	2 00
	(Quinine Sulph. 1-2 gr.)	
	(Morph. Sulph. 1-64 gr.)	
	(Atropine Sul. 1-2000 gr.)	
(Dr. Henry J. Kenyon.)		(Sugar of Milk. 1 gr.)

Four tablets at bedtime, or one every two to four hours—but preferably at bedtime.

Through the vaso-motor nerves, this tablet acts as a specific stimulant to the congested mucous membrane of the naso-pharynx, promotes secretion and favors the return of the diseased parts to their normal condition in from twenty-four to forty-eight hours.

CORYZA No. 4	(Atropine Sulph. 1-600 gr.)	1 15
	(Acid Arsenous. 1-240 gr.)	
	(Strych. Sulph. 1-240 gr.)	
	(Morph. Sulph. 1-60 gr.)	
(Dr. MacCuen Smith.)		(Quinine Sulph. 1-10 gr.)

One or two every hour for four doses, then one every three or four hours if necessary, to control the discharge and headache of acute coryza.

CORYZA No. 5	(Atropine Sulph. 1-600 gr.)	1 15
	(Acid Arsenous. 1-240 gr.)	
	(Strych. Sulph. 1-240 gr.)	
	(Morph. Sulph. 1-100 gr.)	
	(Quinine Sulph. 1-10 gr.)	
(Dr. S. MacCuen Smith.)		(Camphor. 1-4 gr.)

One every hour for cold in the head with weeping nostrils, until the throat feels dry; then one every two or three hours, continuing a day or two after the cold has disappeared.

Please specify Fraser's Tablets

COUGH, ADULTS..... (Dr. Riggs.)	{ Ammon. Mur....3-4 gr. }	2 00
	{ Ext. Opium.....1-15 gr. }	
	{ Fld. Ext. Ipecac. 1 min. }	
	{ Ext. Bellad.....1-30 gr. }	
	{ Ext. Licorice.....1 gr. }	
	{ Oil Anise.....1-20 min. }	

One, every hour or two, for coughs and bronchial irritation.

COUGH, INFANTS..... (Dr. Riggs.)	{ Ammon. Mur....1-4 gr. }	1 25
	{ Tr. Opium.....1-1 min. }	
	{ Fld. Ext. Ipecac. 1-5 min. }	
	{ Ext. Licorice.....1 gr. }	
	{ Tr. Bellad.....1-5 min. }	
	{ Oil Anise.....1-40 gr. }	

Dissolve fifteen tablets in fifteen teaspoonfuls of water and give an infant one teaspoonful every hour or two, for bronchial cough and irritation.

COUGH MIXTURE..... (Dr. Satterlee.)	{ Ammon. Mur....1-4 gr. }	1 00
	{ Paregoric.....5 min. }	
	{ Corrosive Sub....1-6 gr. }	

Give one every hour, or two every two hours during the early stages of cough, and increase the intervals as the symptoms improve.

COUGH No. 2.....	{ Ammon. Carb.....1 gr. }	1 30
	{ Squills Powd.3-4 gr. }	
	{ Senega Powd.....3-4 gr. }	
	{ Paregoric.....5 min. }	

One every hour or two, for persistent cough.

COUGH No. 3.....	{ Tr. Sanguinaria...3 min. }	2 00
	{ Morph. Acet.....1-16 gr. }	
	{ Wine Antimon....5 min. }	
	{ Syr. Wild Cherry....q. s. }	

One every two or three hours, as a sedative, diaphoretic and expectorant.

COUGH, PERSISTENT..... (Dr. Brewer.)	{ Ammon. Mur.....2 gr. }	2 00
	{ Ipecac Powd....1-4 gr. }	
	{ Ext. Squills Acet. 1-4 gr. }	
	{ Fl. Ext. Senega...1 min. }	

One every two or three hours, for cough.

CREOSOTE (Beechwood.).....1-10 gr.....	75
--	----

A tablet three times a day, for diarrhoea and flatulence, and in tuberculosis, gradually increasing the dose.

CREOSOTE (Beechwood.).....1 min.....	1 40
--------------------------------------	------

Give one after meals, for consumption, tubercular diarrhoea, and as an intestinal disinfectant; useful in the abdominal torpidity of epileptics.

CROUP, (Spasmodic.).....	{ Ext. Can. Indica.1-20 gr. }	1 75
	{ Ext. Hyoscyam.1-20 gr. }	
	{ Paregoric.....5 min. }	
	{ Syr. Ipecac.....5 min. }	
	{ Ext. Licorice.....2 gr. }	

Give a tablet every hour until the spasmodic attacks are under control; then diminish the dose.

CROUP No. 2.....	{ Dover's Powder...1-2 gr. }	1 00
	{ Ipecac Powd....1-100 gr. }	
	{ Spongia1-100 gr. }	

One hourly until symptoms are relieved. Give every half hour for three doses, if symptoms are urgent; then one hourly.

When ordering or prescribing

CUBEBS COMP.	{ Rhubarb Powd. 1-2 gr. } { Cubebs Powd. 1-2 gr. } { Calc. Magnesia 1 gr. } { Oil Peppermint 1-20 gr. }	1 10
---------------------------	--	------

A tablet half an hour before meals with a little water, to stimulate the peptic glands and induce appetite.

CYSTITIS	{ Acid Boracic 2 gr. } { Potass. Bicarb. 2 gr. } { Ext. Buchu 1 gr. } { Ext. Triticum Rep. 1 gr. } { Ext. Corn Silk 1-2 gr. } { Ext. Hydrangea 1-2 gr. } { Atropine Sulph. 1-500 gr. }	3 00
-----------------------	--	------

One every three hours, with a large glass of water; for inflammation of the bladder and urinary tract, as well as suppressed renal secretion.

CYSTITIS No. 2	{ Acid Benzoic 1-2 gr. } { Sodium Bicarb. 1-2 gr. } { Oil Gaultheria q. s. }	1 25
-----------------------------	--	------

Two every two hours, generally used for about ten days, for cystitis, gravel and some types of rheumatism.

DAMIANA COMP.	{ Ext. Damiana 2 gr. } { Phosphorus 1-30 gr. } { Ext. Nux Vom. 1-4 gr. }	2 05
----------------------------	--	------

One morning and night for a week, then one every night, as a stimulant to the generative organs.

DAMIANA CO. NO. 2	{ Zinc Phosphide 1-10 gr. } { Ext. Cannab. Ind. 1-4 gr. } { Strych. Sulph. 1-40 gr. } { Ext. Damiana 1-2 gr. }	2 75
--------------------------------	---	------

One after meals, as a nerve tonic.

DAMIANA COMP. No. 3	{ Ext. Damiana 2 gr. } { Ext. Nux Vom. 1-5 gr. } { Cantharides 1-20 gr. }	2 65
----------------------------------	---	------

One tablet three times a day, for impotence.

DIABETIC	{ Croton Chloral 1-2 gr. } { Ext. Opium 1-2 gr. } { Ext. Aloes Barb. 1-6 gr. }	3 00
-----------------------	--	------

One tablet three times a day.

DIARRHŒA	{ Calomel 1-8 gr. } { Morph. Sulph. 1-16 gr. } { Capsicum Powd. 1-16 gr. } { Ipecac Powd. 1-32 gr. } { Camphor 1-16 gr. }	1 15
-----------------------	---	------

One tablet every hour for four doses, then every two hours until relieved.

DIARRHŒA, INFANTS	{ Sodium Salicylate 1 gr. } { Bismuth Subcarb. 1 gr. } { Beta-Naphtol 1-5 gr. }	1 75
--------------------------------	---	------

One every hour or two for fermentative diarrhœa in infants.

DIARRHŒA No. 2	{ Pepsin Pure 1-2 gr. } { Bismuth Subnit. 6 gr. } { Fl. Ext. Nux Vom. 2 min. } { Fl. Ext. Hydrast. 5 min. }	6 25
-----------------------------	--	------

One every two hours for three doses, then every three hours, for diarrhœa caused by indigestion.

Please specify Fraser's Tablets

DIARRHŒA No. 3	{ Opium Powd.....1-4 gr. }	1 15
	{ Camphor.....1-4 gr. }	
	{ Ipecac Powd.....1-8 gr. }	
	{ Lead Acetate.....1-6 gr. }	

Give two at the first dose, then one every hour for six hours, afterwards one after each loose movement, for congestive, irritative and inflammatory diarrhœas.

DIARRHŒA No. 4.....	{ Bismuth Subnit.....2 gr. }	4 25
	{ Salol.....2 gr. }	

(Dr. Tinker.)

One every two hours in acute cases; one every four hours in chronic cases. For gastro intestinal catarrh in adults and children.

DIARRHŒA No. 5.....	{ Bismuth Subnit.....3 gr. }	3 50
	{ Sacch. Pepsin.....2 gr. }	
	{ Arom. Chalk Powd.2 gr. }	

(Board of Health.)

One tablet after each movement; for diarrhœa, accompanied by irritation from undigested food.

DIARRHŒA No. 6.....	{ Bismuth Subnit.....1 gr. }	1 50
	{ Ipecac Powd.....1-20 gr. }	
	{ Opium Powd.....1-40 gr. }	

(Dr. Elliott Riggs.)

For Children.

One every two or three hours to a child about one year old, as may be necessary to check diarrhœa; preceded by eight to ten doses Bismuth and Ipecac Comp. No. 2, taken one every three hours.

DIARRHŒA No. 7	{ Corros. Sublim....1-96 gr. }	3 75
	{ Bismuth Subnit.....5 gr. }	
	{ Oil Anise.....q. s. }	

(Dr. Bierhoff.)

One, every one or two hours, for summer diarrhœa in children, where an antiseptic effect is desired.

DIARRHŒA No. 8.....	{ Opium Powd.....1-8 gr. }	1 00
	{ Ipecac Powd.....1-15 gr. }	
	{ Lead Acetate.....1-4 gr. }	

(Dr. Drumheller.)

One tablet every two or three hours, for diarrhœa.

DIARRHŒA No. 9.....	{ Comp. Chalk Powd.1 gr. }	1 00
	{ Acid Tannic.....1-20 gr. }	
	{ Opium Powd.....1-40 gr. }	
	{ Camphor.....1-40 gr. }	

One crushed and taken on the tongue or diffused in water on the end of a spoon, every twenty minutes, for a baby with diarrhœa, pain or flatulence; after the sixth dose, give every hour.

DIARRHŒA No. 10.....	{ Camphor Water...4 min. }	3 75
	{ Paregoric4 min. }	
	{ Bismuth Subnit....4 gr. }	
	{ Svr. Rhub. Arom.20 min. }	
	{ Oil Peppermint.1-8 min. }	

(Dr. Foster.)

Two tablets every one, two or three hours, as case demands.

DIARRHŒA No. 11.....	{ Salol.....1-2 gr. }	2 75
	{ Bismuth Salicylate...1 gr. }	
	{ Calomel.....1-40 gr. }	
	{ Paregoric.....2 min. }	
	{ Mist. Cret. Arom....q. s. }	
	{ Oil Cinnamon.....q. s. }	

(Dr. Prendergast.)

One or two tablets every hour until patient is better; for diarrhœa, accompanied by colicky pains, and where an intestinal antiseptic is indicated.

DIFFUSIVE MALARIAL.....	{ Corrosive Sub....1-50 gr. }	1 25
	{ Ammon. Mur.....2 gr. }	
	{ Cinchonia Sulph.1-10 gr. }	
	{ Cinchonid. Sulph.1-10 gr. }	

{ Quinine Sulph....1-10 gr. }

One to three tablets after meals, preceded by three one grain calomel tablets at intervals of an hour; for dumb ague, headache and cachexia of malarial origin.

DIGESTIVE.....	<div> <div>Pepsin.....2 gr.</div> <div>Pancreatin.....1 gr.</div> <div>Ipecac Powd.....1-10 gr.</div> <div>Jamaica Ginger...4-5 gr.</div> <div>Bismuth Subnit.....2 gr.</div> <div>Ext. Nux Vom...1-10 gr.</div> <div>Oil Wintergreen....q. s.</div> </div>	6 75
----------------	---	------

One or two after meals, repeated in an hour, for cases of impaired or slow digestion requiring a stomachic and carminative in addition to the ordinary solvents for food.

DIGESTIVE TONIC.....	<div> <div>Iron Lactate.....1 gr.</div> <div>Quinine Sulph.....1 gr.</div> <div>Acid Arsenous...1-30 gr.</div> <div>Strych. Sulph....1-30 gr.</div> <div>Lactopeptine.....2 gr.</div> </div>	
(Dr. Satterlee.)	Per 100.....	1 25

One tablet before each meal, as a general tonic and appetizer, as well as digestive. Is well borne by the most delicate stomach.

DIGITALIA, Merck's.....	1-1000 gr.....	75
DIGITALIA, Merck's.....	1-500 gr.....	75
DIGITALIA, Merck's.....	1-400 gr.....	1 00
DIGITALIA, Merck's.....	1-300 gr.....	1 10
DIGITALIA, Merck's.....	1-200 gr.....	1 30
DIGITALIA, Merck's.....	1-150 gr.....	1 70
DIGITALIA, Merck's.....	1-100 gr.....	2 25
DIGITALIA, Merck's.....	1-60 gr.....	2 75
DIGITALIA, Merck's.....	1-50 gr.....	3 00
DIGITALIS COMP.....	<div>Tr. Digitalis.....2 min.</div> <div>Brucia.....1-100 gr.</div>	1 00
(Dr. Wood.)		

Give a tablet every three hours, as a heart stimulant.

DIGITALIS COMP. No. 2.....	<div>Tr. Digitalis... 2 min.</div> <div>Strych. Sulph... 1-100 gr.</div>	1 00
(Dr. Wood.)		

One every three hours, as a heart stimulant and tonic.

DIGITALIS POWD.....	1-20 gr.....	75
Heart tonic and diuretic.		

One every three hours for a child, to regulate the heart and increase the flow of urine.

DIGITALIS POWD.....	1-10 gr.....	75
---------------------	--------------	----

A tablet every three hours, for a child as a diuretic and to strengthen cardiac systole.

DIGITALIS POWD.....	1-5 gr.....	75
---------------------	-------------	----

Give one every three hours to assist absorption of serous exudations and as a heart tonic.

DIGITALIS POWD.....	1-4 gr.....	75
---------------------	-------------	----

One every three or four hours, as a heart tonic.

DIGITALIS POWD.....	1-2 gr.....	75
---------------------	-------------	----

One tablet three times a day, as a renal and cardiac tonic.

DIGITALIS FL. EXT.....	1-2 min.....	75
------------------------	--------------	----

One every three or four hours, for mitral disease, partial suppression of urine, and dropsy.

DIGITALIS FL. EXT.....	1 min.....	1 00
------------------------	------------	------

Give one after meals to control excessive and irregular heart action and stimulate the kidneys; reduce the dose if headache and flushing occurs.

Please specify Fraser's Tablets

DIGITALIS & IRON COMP.	Ext. Digitalis....1-10 gr.	1 00
Cardiac tonic and haematic.	Iron Phosphate....1-4 gr.	
	Strych. Sulph....1-60 gr.	

A tablet after meals, for anæmia with palpitation of the heart, and in convalescence from exhausting illness.

DIPHTHERIA AND GRIP.	Hydrarg. Bichlor 1-50 gr.	2 35
(Dr. Briggs.)	Guaiaac1 gr.	
	Tr. Sanguinaria...3 min.	
	Tr. Colchicum.....6 min.	
	Tolu1-5 gr.	
	Powd. Rhei.....1-2 gr.	
	Cubebs Pulv.....1 1-5 gr.	

Dissolve a tablet slowly in the mouth every two or three hours, supplement with a heart tonic and quinine, also with a salicylate of sodium tablet if needed for pain.

DIPSOMANIA	Gold Chloride...1-40 gr.	2 00
	Strych. Nitrate...1-60 gr.	

One tablet every two or three hours, reducing to three a day, to control the craving for alcohol and as a tonic after a debauch.

DIURETIC	Digitalis Powd....1 gr.	1 75
	Ext. Buchu.....1 gr.	
	Potass. Nitrate...1 gr.	
	Squills Powd.....1 gr.	

One, three times a day, followed by a large glass of water.

DIURETIC No. 3	Strych. Sulph....1-100 gr.	2 50
	Caffeine Hydrob....1 gr.	
	Digitalis Powd....1-2 gr.	

One to three tablets three times a day, as a mild, non-irritating diuretic

DIURETIC No. 4	Digitalis Powd....1 gr.	2 50
(Dr. Christy.)	Potass. Nitrate....2 gr.	
	Ext. Buchu1-2 gr.	
	Ext. Scoparius....1-2 gr.	
	Oil Juniper.....1 gtt.	

One every two hours as a diuretic; after the third dose give every four hours, for suppression of urine, dropsy and chronic Bright's disease; omit the medicine one or two days each week.

DOVER'S COMP.	Quinine Sulph....1-2 gr.	1 50
(Dr. Terriberry.)	Dover's Powd....1-2 gr.	

To an infant under six months, one every two hours at first, in syrup; over six months and up to two years, give one every hour at first, then diminish the dose as the effect is felt. Used in acute bronchitis and pneumonia of children.

DOVER'S COMP. No. 2	Dover's Tablets 2 1-2 gr.	1 35
	Calomel1-4 gr.	

One every hour for three doses, with a large glass of water, for colds with fever.

DOVER'S COMP. No. 3	Dover's Powder1 gr.	1 25
	Calomel1-2 gr.	

Give one tablet every three hours as necessary.

DOVER'S POWDER	1-10 gr.	75
-----------------------------	----------	----

Dissolve ten in four tablespoonfuls of water; of this give a teaspoonful every half hour, with a half grain blue mass tablet every hour, for infantile diarrhœa with offensive passages.

DOVER'S POWDER	1-5 gr.	75
-----------------------------	---------	----

One every half hour for five doses, for a child with fever or bronchitis.

DOVER'S POWDER	1-4 gr	75
-----------------------------	--------	----

One tablet every half hour to four doses for a child, to abort a cold after exposure; useful also as a sudorific in fevers of childhood.

When ordering or prescribing

DYSPEPTIC, SENILE	{ Aloin.....1-20 gr. }	1 25
(Dr. Hawes.).....	{ Pepsin, Pure.....1-5 gr. }	
	{ Strych. Salph....1-100 gr. }	

A tablet before and after meals, for the slow digestion and mal-assimilation of old age.

EISENZUCKER	3 grs.....	1 00
--------------------------	------------	------

Children's iron tonic.

One to three tablets, three times a day after meals, as an iron tonic for children.

EISENZUCKER	5 grs.....	1 25
--------------------------	------------	------

One to three tablets three times a day after meals, as an iron tonic.

ELATERIN CRYSTAL	1-10 gr.....	7 50
-------------------------------	--------------	------

Purgative.

One every three hours for three doses if necessary, where a strong purgative effect is required; best combined with inspissated ox gall, about three grains to each dose of elaterin.

ELATERIUM, Clutterbuck's	1-10 gr.....	2 00
---------------------------------------	--------------	------

One at night as a purge; one, repeated in an hour, followed by a saline, for suppression of urine and where sure action is necessary.

EMMENAGOGUE	{ Ergotin.....1 gr. }	4 00
	{ Ext. Gossypium....1 gr. }	
	{ Iron Sulph Dry....1 gr. }	
	{ Ext. Bl'k Hellebore.1 gr. }	
	{ Aloes Soc.....1 gr. }	
	{ Oil Savine.....1-4 min. }	

One every two hours as an emmenagogue.

EMMENAGOGUE No. 2	{ Pulv. Aloes Soc....1 gr. }	1 65
	{ Ferri Sulph. Exsic.1-2 gr. }	
	{ Ext. Helleb. Nig..1-4 gr. }	
	{ Pulv. Myrrh.....1-4 gr. }	
	{ Pulv. Sapo.....1-4 gr. }	
	{ Pulv. Canella....1-8 gr. }	
	{ Pulv. Zingib. Jam..1-8 gr. }	

One to three tablets three times daily.

ERGOTIN	1-50 gr.....	75
----------------------	--------------	----

One every two hours to a child, as a general nerve tonic and vaso-motor stimulant.

ERGOTIN	1-20 gr.....	75
----------------------	--------------	----

A tablet every three hours to a child, as a general nerve tonic and vaso-motor stimulant.

ERGOTIN	1-10 gr.....	75
----------------------	--------------	----

One three times a day for a general vaso-motor tonic in chorea and epilepsy of children.

ERGOTIN	1-6 gr.....	95
----------------------	-------------	----

One three times a day as a general nerve tonic and to counteract tendency to internal hemorrhage.

ERGOTIN	1-4 gr.....	1 15
----------------------	-------------	------

One to four, repeated as required for nerve tonic.

ERGOTIN	1-2 gr.....	1 70
----------------------	-------------	------

One every hour or two as an oxytocic and to control internal bleeding.

ERGOTIN	1 gr.....	2 75
----------------------	-----------	------

A tablet every hour, as necessary, to control hæmoptysis, and hæmorrhage from the bowel; also after parturition to secure uterine contraction.

ERGOTIN	2 gr.....	4 25
----------------------	-----------	------

One every three hours to control internal hemorrhage, and as a general nerve tonic.

ERGOTIN COMP.	{ Ergotin.....1 1-2 gr. }	3 00
General tonic.	{ Tr. Digitalis.....1-2 min. }	
	{ Quinine Sulph.....1-2 gr. }	

One to three tablets every three hours for nerve and heart tonic.

ERGOTIN COMP. No. 2	{ Ergotin.....1-2 gr. }	2 25
	{ Ext. Can. Ind.....1-4 gr. }	

One every three hours.

EPILEPTIC	{ Sodium Bromide....3 gr. }	3 00
(Dr. Batton.)	{ Ammon. Bromide....3 gr. }	
	{ Potass. Bromide....3 gr. }	
	{ Tr. Calab. Bean 2 1-2 min. }	
	{ Ext. Hyoseyam....1-2 gr. }	

Two tablets three times daily with a swallow of water, may be given for a year if required, for epilepsy.

EPILEPTIC No. 2	{ Sodium Bromide....3 gr. }	2 75
(Dr. Batton.)	{ Ammon. Bromide....3 gr. }	
	{ Potass. Bromide....3 gr. }	
	{ Tr. Calab. Bean 2 1-2 min. }	

Two tablets three times daily with a swallow of water; may be used continuously for a long period.

ESERINE SULPHATE1-100 gr.....	2 50
--	------

Cerebral tonic.

A tablet three times a day as a cerebral tonic in tetanoid and choreic conditions.

ESERINE SULPHATE1-50 gr.....	1 75
---	------

A tablet one to three times a day in convulsive affections and intestinal atony.

EUONYMIN1-10 gr.....	1 00
-----------------------------------	------

Hepatic.

One every hour for five doses, in hepatic torpor.

EUONYMIN1-8 gr.....	1 15
----------------------------------	------

A tablet every hour for three doses, followed by an aperient, for torpidity of the liver.

EUONYMIN1-4 gr.....	1 50
----------------------------------	------

One three times a day, with half an ounce of Castor Oil at night, for torpidity of the liver and duodenal catarrh.

EUONYMIN1-2 gr.....	2 25
----------------------------------	------

One to three a day, in divided doses, followed by an aperient, for deficiency of bile in the intestines.

EUONYMIN COMP.	{ Euonymin.....1-4 gr. }	1 70
(Dr. Welch.)	{ Podophyllin.....1-8 gr. }	
	{ Aloin.....1-8 gr. }	

One every night for a week, and one after meals every second day, if necessary, in cases of hepatic torpor with biliousness, headache and light colored fæces.

EXALGINE2 grs.....Per 100	1 50
--	------

Antipyretic.

A tablet every three hours to relieve the pain of rheumatism and neuralgia.

EXALGINE3 grs.....Per 100	2 25
--	------

One every three hours for neuralgia and rheumatic pains; one three times a day for diabetes, to lessen the quantity of urine and control the sugar excretion.

EXALGINE5 grs.....Per 100	3 25
--	------

One or two a day for chorea; also a single dose for neuralgia and rheumatic pain.

EXPECTORANT (Dr. Rankin.)	Bellad. Fl. Ext.1-8 min.	2 25
	Ipecac Powd.1-10 gr.	
	Ext. Licorice.1-8 gr.	
	Codeine1-16 gr.	
	Ext. Senega1-10 gr.	

One every hour or two, for an irritative bronchial cough.

EXPECTORANT No. 4 (Dr. Bierhoff.)	Ammon. Carb.1 gr.	1 25
	Ammon. Muriate.1 gr.	
	Ext. Licorice1-2 gr.	
	Powd. Squill1-4 gr.	
	Oil Cinnamon.....q. s.	

One every hour or two, as a stimulant and expectorant in laryngitis and bronchitis of children; best given dissolved in one or two teaspoonfuls of water.

EXPECTORANT, Anodyne No. 2 (Dr. Bolton.)	Ammon. Mur.1 gr.	1 75
	Tartar Emetic.1-30 gr.	
	Sanguinar. Nit.1-100 gr.	
	Morphine Sulph.1-50 gr.	

One or two tablets every two hours, as an anodyne in coughs with respiratory inflammation.

FEBRIFUGE	Quinine Sulph.2 gr.	1 00
	Acetanilid.2 gr.	
	Cocaine Muriate.1-6 gr.	
	Per 100	

One every two hours, with free use of drinking water, for fevers of the continued type.

FEVER No. 2 (Dr. Davis.)	Tr. Aconite.1-5 min.	1 00
	Tr. Bryonia.1-10 min.	
	Tr. Belladonna.1-10 min.	

One tablet every two or three hours, with plenty of water, for the fevers and ordinary colds of infancy and childhood. Give adults two to four tablets at a dose.

FEVER, HECTIC (Dr. Westbrook.)	Acid Arsen.1-20 gr.	4 00
	Ext. Nux Vom.1-3 gr.	
	Quinine Sulph.2 gr.	
	Acetanilid.3 gr.	

Give three to four tablets before four o'clock in the afternoon, to prevent occurrence of hectic fever.

FEVER, Improved. (Dr. Kenyon.)	Tr. Aconite.1 min.	1 50
	Morph. Sulph.1-50 gr.	
	Tartar Emetic.1-50 gr.	
	Ipecac Powd.1-4 gr.	

One every hour for six doses, then every two to four hours; for cold with fever, to relieve the sensation of tightness or fulness in the bronchial tubes, and in the early stage of bronchitis.

FEVER, INFANTS	Aconite.1-100 gr.	75
	Ipecac Powd.1-100 gr.	

One every hour for ten doses, afterwards at intervals of two to four hours.

FEVER & AGUE	Copper Sulph.1-10 gr.	2 00
	Opium Powd.1-8 gr.	
	Quinine Sulph.1-2 gr.	
	Podophyllin1-12 gr.	

One to six a day for fevers due to malarial infections; occasionally as a preventive in malarious districts.

FOUR CHLORIDES Tonic.	Iron Chloride.1-2 gr.	3 00
	Arsenic Chlor.1-64 gr.	
	Merc. Bichlor.1-48 gr.	
	Quinine Chloride.1 gr.	

One just after each meal for tonic in anæmia, and as an alterative.

FOUR CHLORIDES No. 2	{ Iron Chloride.....1-2 gr. }	2 00
	{ Arsenic Chlor.1-100 gr. }	
	{ Merc. Bichlor.1-100 gr. }	
	{ Quinine Chloride.1-2 gr. }	

One after each meal. A useful tonic in all kinds of anæmia.

GALLIC ACID COMP.	{ Gallic Acid.....3 gr. }	1 85
	{ Ext. Belladonna.1-5 gr. }	

General astringent.

One tablet as necessary for hæmatemesis, menorrhagia and as a general astringent.

GARGLE	{ Sodium Bicarb.3 1-3 gr. }	2 60
	{ Sodium Salicyl.3 1-3 gr. }	
	{ Acid Boracic.3 1-3 gr. }	
	{ Oil Wintergreen.q. s. }	

Three tablets to a cupful of hot water; gargle every hour until relieved, using it hot if possible.

GENTIAN, ALKALINE	{ Infus. Gent. Co.1 teasp. }	1 25
	{ Sodium Bicarb.5 gr. }	

(Dr. J. J. Moran.)

Two or three tablets dissolved in water before each meal. A tonic stomachic where there is catarrhal gastric disorder, and ordinary atonic dyspepsia.

GELSEMIN	1-16 gr.....	1 25
-----------------------	--------------	------

Antispasmodic and relaxant.

One every hour for three doses, for neuralgia, toothache and nervous cough.

GELSEMIN	1-10 gr.....	1 50
-----------------------	--------------	------

A tablet, repeated in an hour, together with hot baths, for dental neuralgia; and for hepatic and renal colic,

GELSEMIN	1-8 gr.....	1 65
-----------------------	-------------	------

A tablet three times a day as a depressant in pleurisy and pneumonia, and for cough with a strong nervous element.

GELSEMIN	1-4 gr.....	2 35
-----------------------	-------------	------

One or two doses at intervals of two hours, for dental neuralgia, headache, and to control the paroxysms of asthma.

GELSEMINE SULPH	1-100 gr.....	3 75
------------------------------	---------------	------

Three to five a day in divided doses, for spermatic and neuralgic affections and as a sedative to the spinal cord.

GELSEMINE SULPH	1-50 gr.....	6 75
------------------------------	--------------	------

A tablet every three or four hours, for neuralgia; also as an antidote to Strychnine.

GELSEMIUM FL. EXT.	1 min.....	1 00
---------------------------------	------------	------

One every hour to six doses, in the onset of fevers, especially of malarial origin, and for various neuralgias.

GOLD & SODIUM CHLORIDE	1-30 gr.....	1 30
---	--------------	------

Tonic.

A tablet after meals as a general tonic, for anæmia and after wasting disease.

GOLD & SODIUM CHLORIDE	1-20 gr.....	1 70
---	--------------	------

A tablet after meals as a general tonic, for anæmia and after wasting disease.

GOLD & SODIUM CHLORIDE	1-10 gr.....	3 00
---	--------------	------

One with meals, and one at night, to combat alcohol and morphine craving, to hasten the elimination of syphilitic poison, and as a hæmal tonic.

GONORRHOEA	{ Cubebs Pulv.1 gr. }	2 25
	{ Copaiba Solid.1 gr. }	
	{ Iron Sulph. Dry.1-4 gr. }	
	{ Oil Sandal.1-1 min. }	
	{ Oil Wintergr.1-10 min. }	
	{ Venice Turpen.1-4 min. }	

Two tablets, three or four times daily, usually supplemented with ten grains of bicarbonate of sodium or potassium.

GONORRHŒA No. 2	{ Cubebs.....1 1-4 gr. }	
(Dr. Burd.)	{ Copaiba Solid...1 1-4 gr. }	1 50
	{ Iron Sulph. Dry...1-4 gr. }	
	{ Venice Turpent...1-2 gr. }	

One tablet every two hours, or two tablets every three hours; used in catarrhal or subacute stage of gonorrhœa.

GRAY POWDER COMP	{ Merc. with Chalk...1 gr. }	
(Dr. Anderton.)	{ Salol.....1 gr. }	3 75
	{ Bismuth Subnit.....3 gr. }	

One every hour until ten doses have been taken; for torpid liver, loose, colorless, mal-odorous stools, with or without vomiting.

GRAY POWDER CO. No. 2 ...	{ Merc. with Chalk...1 gr. }	
(Dr. Hutchinson.)	{ Dover's Powder....1 gr. }	1 35

Give one tablet every hour or two until symptoms improve; then every three hours, for a child with cough, fever and constipation.

GRIP	{ Ammon. Salicyl....3 gr. }	
	{ Phenacetine.....1 gr. }	8 00
	{ Caffeine.....1-2 gr. }	

A tablet every two or three hours in the febrile stages of influenza, until the cardiac depression and the fever improve.

GRIP No. 3	{ Phenacetine.....1 gr. }	
(Dr. Searles.)	{ Salicin.....1 gr. }	7 00
	{ Quinine Sulph....1-2 gr. }	

One, cautiously, preceded by a cathartic, every hour for three doses, then one every two hours, for grip.

GUAIAC COMP	{ Guaiac.....1 2-3 gr. }	
Throat.	{ Ammon. Mur....1 2-3 gr. }	1 10
	{ Ext. Licorice....1-2 gr. }	

One slowly dissolved on the tongue, for astrigent and antiseptic effects upon the throat.

HÆMATIC	{ Acid Arsen.....1-120 gr. }	
(Dr. Clymer.)	{ Iron Powd.....1-10 gr. }	75
	{ Merc. Sub. Cor..1-200 gr. }	
	{ Nux Vomica. Po..1-20 gr. }	

Give one about an hour after meals, as a hæmatic tonic in anæmia.

HÆMOGALLOL	3 grs.....	Per 100	1 75
-------------------------	------------	---------	------

Begin with one after meals; the second week, give two at a dose; if well tolerated, give three for the third week and after; for anæmia, chlorosis and menstrual disorder dependent upon them.

HEADACHE	{ Acetanilid.....1 gr. }	
(Dr. Chamberlain.)	{ Ext. Belladonna...1-8 gr. }	1 15
	{ Ext. Gelsemium...1-4 gr. }	

One or two at first and repeat every two to four hours, as required; for congestion or neuralgic headache, with fever.

HEART STIMULANT	{ Tr. Digitalis.....2 min. }	
	{ Tr. Strophanthus..2 min. }	1 75
	{ Tr. Belladonna...1-4 min. }	
	{ Nitroglycerin....1-100 gr. }	

One every three hours for smokers heart, palpitation, and angina pectoris.

HEART STIMULANT No. 2	{ Tr. Digitalis.....2 min. }	
	{ Tr. Strophanthus..2 min. }	1 75
	{ Tr. Belladonna...1-4 min. }	
	{ Nitroglycerin....1-200 gr. }	

One every three hours for smokers heart, palpitation, and angina pectoris.

HEART STIMULANT No. 3	{ Atropine Sulph....1-75 gr. }	
(Dr. B. F. Horner.)	{ Strych. Sulph....1-30 gr. }	1 25

One tablet every three hours for certain cases of slow heart, attended with valvular lesions at aortic or pulmonary orifices, and of some particular forms at mitral valve.

HEART STIMULANT No. 4.... (Dr. Durant.)	{ Fl. Ext. Digitalis... 1 min. Nitroglycerin... 1-100 gr. Strych. Sulph.... 1-50 gr. }	1 35
---	--	------

One tablet, three or four times a day, when digitalis is indicated in chronic valvular diseases.

HEART TONIC.....	{ Fl. Ext. Cac. Grand. 5 min. Tinct. Digitalis... 2 min. Caffeine Cit..... 2 gr. }	5 05
-------------------------	--	------

One every three hours for weak heart or feeble heart of aged.

HEART TONIC No. 2..... (Dr. Gordinier.)	{ Nitroglycerin... 1-200 gr. Tr. Strophanthus... 3 min. Fl. Ext. Digitalis... 1 min. Strych. Sulph.... 1-60 gr. Iron Powd..... 1 gr. }	2 00
---	--	------

Give one three times daily, or if occasion requires, every three or four hours, as a general heart tonic, and tonic in anæmia.

HEART TONIC No. 3..... (Dr. Pultz.)	{ Tr. Digitalis..... 1 min. Tr. Strophanthus... 1 min. Tr. Cactus Grand... 1 min. Sparteine Sulph. 1-100 gr. Nitroglycerin... 1-100 gr. }	1 50
---	---	------

One tablet every three hours for irregular and weak heart.

HELLEBOREIN..... Cardiac tonic.	1-100 gr.....	1 50
---	---------------	------

One every two hours, until the pulse is under control, then one every four to six hours, as a heart tonic where digitalis disagrees.

HELLEBOREIN.....	1-50 gr.....	2 60
-------------------------	--------------	------

A tablet two or three times a day, to strengthen cardiac systole and as a general vaso-motor stimulant; increase with caution.

HELONIN..... Diuretic and tonic.	1-10 gr.....	1 30
--	--------------	------

One three times a day to increase the flow of urine, and as an emmenagogue.

HEMICRANIA.....	{ Croton Chloral..... 1 gr. Ext. Cannab. Ind... 1-4 gr. }	3 50
------------------------	--	------

Give one tablet and another in half an hour; for nervous or sick headache.

HEPATIC..... (Dr. Henry J. Kenyon.)	{ Euonymin... 1-8 gr. Podophyllin... 1-20 gr. Ipecac Powd..... 1-8 gr. Calomel..... 1-8 gr. Aloin..... 1-12 gr. }	1 15
---	---	------

HEPATIC ALTERNATIVE..... (Dr. Temple.)	{ Bismuth Submt..... 1 gr. Salol..... 1 gr. Calomel..... 1-10 gr. }	2 50
--	---	------

One tablet every hour until laxative effect is produced, then two or three times a day for a week or more; for bilious dyspepsia with flatulence.

HYDRARG. COMP. No. 1..... (Dr. Leaming.) Bronchial expectorant.	{ Calomel..... 1-24 gr. Tartar Emetic... 1-120 gr. Nitr. Potass.... 2 1-2 gr. Ext. Licorice... .. Oil Sassafras..... Oil Wintergreen..aa. q.s. }	1 10
--	---	------

One every three hours for cough after grip.

Please specify Fraser's Tablets

HYDRARG. COMP. No. 2 (Dr. Leaming.)	{ Calomel.....1-12 gr. }	1 50
	{ Tartar Emetic...1-60 gr. }	
	{ Nitr. Potass.....5 gr. }	
	{ Ext. Licorice..... }	
	{ Oil Sassafras..... }	
	{ Oil Wintergreen, aa. q. s. }	

Give a tablet every four hours as an expectorant in bronchitis and influenza.

HYDRARG. COMP. No. 3 (Dr. Leaming.)	{ Calomel.....1-6 gr. }	2 00
	{ Tartar Emetic...1-30 gr. }	
	{ Nitr. Potass.....10 gr. }	
	{ Ext. Licorice..... }	
	{ Oil Sassafras..... }	
	{ Oil Wintergreen, aa. q. s. }	

One after meals, for chronic inflammation of the bronchial tubes with thick and scanty sputa, bronchial dilatation and emphysema.

HYDRARG. & IPECAC COMP. Alterative and hepatic.	{ Mass Hydrarg.....1 gr. }	1 50
	{ Ipecac Powd.....1-8 gr. }	
	{ Opium Powd.....1-4 gr. }	

One after meals with an occasional saline in the morning, for loss of function in the liver, with biliousness, headache and constipation.

HYDRARG. & PODOPHYLLIN Hepatic.	{ Mass Hydrarg.....1 gr. }	1 10
	{ Podophyllin.....1-4 gr. }	

One to four at bedtime, as a hepatic laxative in duodenal catarrh and intestinal dyspepsia.

HYDRASTIA, (White Alkaloid.) Stimulant.	1-20 gr.....	1 10
--	--------------	------

One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.

HYDRASTIN.....	1-100 gr.....	75
----------------	---------------	----

One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.

HYDRASTIN.....	1-20 gr.....	1 10
----------------	--------------	------

One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.

HYDRASTIN MUR.....	1-100 gr.....	75
--------------------	---------------	----

One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.

HYDRASTIN CAPSICUM CO.. (Dr. Postle.)	{ Hydrastin.....1-4 gr. }	2 00
	{ Podophyllin.....1-10 gr. }	
	{ Capsicum Powd...1-4 gr. }	

One to three tablets a day, between meals, as a stomachic tonic and laxative, and also as an appetizer.

HYDRASTIN COMP.....	{ Podophyllin.....1-20 gr. }	1 25
	{ Hydrastin.....1-10 gr. }	

One every three hours, for not more than three doses, to regulate the intestines in atonic conditions, decreasing to one or two at night.

HYOSCIN HYDROBROM.....	1-200 gr.....	3 40
------------------------	---------------	------

Anodyne.

One every hour or two, as necessary to quiet the nerves and induce sleep, and to supplement the action of other hypnotics.

HYOSCIN HYDROBROM.....	1-100 gr.....	6 00
------------------------	---------------	------

One tablet every hour for three doses, to control the paroxysms of asthma and for neuralgia. Must be given with caution in the case of a weak heart.

HYOSCIN HYDROBROM.....	1-50 gr.....	11 00
------------------------	--------------	-------

One as a nerve sedative, for mania, various neurotic conditions, asthma, and as an anodyne in pain; repeated with caution.

HYOSCYAMIA, (White Cryst.)	1-500 gr.....	1 50
A tablet every two hours, for cough with pain and restlessness.		
HYOSCYAMIA, (White Cryst.)	1-400 gr.....	1 85
One tablet repeated in an hour, as a hypnotic and sedative.		
HYOSCYAMIA, (White Cryst.)	1-300 gr.....	2 00
HYOSCYAMIA, (White Cryst.)	1-200 gr.....	3 40
HYOSCYAMIA, (White Cryst.)	1-100 gr.....	6 00
HYOSCYAMIA, (White Cryst.)	1-50 gr.....	11 00
HYOSCYAMUS COUGH	{ Codeine.....1-10 gr. { Ext. Hyoscyam...1-8 gr. }	2 25
One every three hours in cough accompanied by pain or a strong nervous element.		
HYOSCYAMUS EXT.	1-4 gr.....	75
Sedative.		
One tablet every four hours, for vesical tenesmus and dysuria, and cough with pain.		
HYOSCYAMUS EXT.	1-2 gr.....	1 15
A tablet three times a day to control the pain of inflammation in the respiratory and uro-poietic systems and as a general nerve sedative.		
HYPNOTIC	{ Sodium Bromide,...5 gr. Acetanilid.....2 gr. Hyoscyamia....1-400 gr. Digitalia.....1-400 gr. }	3 50
One every two hours to produce sleep in threatened delirium and mania, and for the pain of various inflammations.		
HYPOPHOSPHITES & QUININE	{ Quinine Hypo.....1 gr. Ferrum Hypo....1-2 gr. Calcium Hypo....1-2 gr. Sodium Hypo....1-4 gr. Potass. Hypo....1-4 gr. Mangan. Hypo....1-4 gr. Strych. Hypo....1-64 gr. }	3 50
Tissue tonic.		
A tablet after meals for anæmia, and in convalescence from wasting disease.		
HYPOPHOSPHITES & QUININE	{ Quinine Hypo.....1-2 gr. Ferrum Hypo....1-4 gr. Calcium Hypo....1-4 gr. Sodium Hypo....1-8 gr. Potass. Hypo....1-8 gr. Mangan. Hypo....1-8 gr. Strych. Hypo....1-128 gr. }	2 25
1-2 Strength.		
For a child, one after meals, for anæmia and loss of flesh.		
HYPOPHOSPHITES & QUININE WITH CREASOTE	{ Quinine Hypo.....1 gr. Ferrum Hypo....1-2 gr. Calcium Hypo....1-2 gr. Sodium Hypo....1-4 gr. Potass. Hypo....1-4 gr. Mangan. Hypo....1-4 gr. Strych. Hypo....1-64 gr. Creasote.....1-2 min. }	3 75
One after meals, for beginning phthisis, to prevent emaciation and loss of strength, or as a general chemical food and intestinal antiseptic.		
IGNATIA POWD.	1-100 gr.....	75
Tonic.		
One to a child every two hours as a stomachic and general tonic.		
IGNATIA POWD.	1-20 gr.....	75
One tablet every two hours, as a gastric stimulant and nerve tonic.		

IGNATIA POWD......1-10 gr. 75

One tablet every two hours, as a gastric stimulant and nerve tonic.

IGNATIA COMP...... } Ignatia Powd...1-100 gr. { 75
(Dr. Gilbert.) } Ipecac Powd....1-10 gr. {

One an hour after meals, for nervous dyspepsia.

INDIGESTION...... } Nux Vomica....1-50 gr. { 75
(Dr. Gilbert.) } Carbo Veg.....q. s. {

One tablet every two hours for indigestion from excessive eating or drinking.

INDIGESTION No. 3...... } Alpha Naphtol...2 1-2 gr. { 5 00
 } Bism. Subcarb...1 1-2 gr. {
 } Rhubarb Powd....1 gr. {
 } Codeine1-20 gr. {
 } Ext. Bellad....1-20 gr. {
 } Ext. Nux Vom...1-20 gr. {

Two tablets immediately after each meal followed by one an hour after, and two more at bedtime; for intestinal indigestion.

INDIGESTION POWD. No. 1 } Pepsin Sacch....1-40 gr. { 75
 } Carbo Veg.....1-400 gr. {
 } Bismuth Subnit..1-40 gr. {

A tablet crushed in a teaspoonful of sweetened water, for an infant with dyspepsia and vomiting; give fifteen minutes before and directly after feeding.

INDIGESTION POWD. No. 2. } Carbo Veg.....1-400 gr. { 75
 } Rhubarb Powd..1-40 gr. {
 } Bismuth Subnit..1-40 gr. {
 } Pepsin.....1-40 gr. {

One after feeding, for the fermentative and acid dyspepsia of nursing infants; crush the tablet in a teaspoonful of water, which may be sweetened if desired.

INFLAMMATUS...... } Tr. Verat Vir...1-12 min. { 1 75
(Dr. Seward.) } (Norwood's)
 } Tr. Aconite Rt..1-12 min. {
 } Tr. Opium.....2-3 min. {
 } Tr. Bryonia2-3 min. {
 } Tr. Belladonna..2-3 min. {
 } Tr. Cimicifuga..2-3 min. {
 } Tr. Guaiac2-3 min. {
 } Tr. Colch. Seed..2-3 min. {
 } Tr. Colocynth...2-3 min. {

One every two to four hours, or one dissolved in fifteen teaspoonfuls of water, of this a teaspoonful every fifteen minutes; for acute stages of rheumatism or neuralgia from exposure, also for early stages of tonsillitis.

INFUSION GENTIAN COMP...... 1 25

Each tablet represents one drachm Compound Infusion Gentian, U. S. P., 1870.

From two to four tablets after each meal as a stomachic and carminative.

INJECTION..... } Hydrastin.....1-5 gr. { 2 75
(Dr. Briggs.) } Zinc Sulph.....3 gr. {
 } Morph. Sulph....1-20 gr. {
 } Atropine Sulph...1-50 gr. {

Dissolve one tablet in half a glass of warm water and make three injections carefully after each urination; warm the solution for each using, for best results.

IPECAC POWD......1-1000 gr. 75

One every three hours for an infant, as a gentle stimulant to the gastric and intestinal mucous membrane.

IPECAC POWD.	1-200 gr.	75
One every three hours, for the atonic dyspepsia of anæmic children.		
IPECAC POWD.	1-100 gr.	75
A tablet three or four times a day, for cough and bronchitis in children, as a stimulating expectorant.		
IPECAC POWD.	1-50 gr.	75
One every hour, for diarrhœa and dysentery in children.		
IPECAC POWD.	1-10 gr.	75
One every three hours for a child, as a vascular and hepatic stimulant; every hour as a sudorific and expectorant in diseases of the respiratory organs.		
IPECAC POWD.	1-4 gr.	90
Emetic and expectorant.		
As an emetic for an infant, one repeated if necessary; one every four hours as an expectorant in cough.		
IPECAC POWD.	1-2 gr.	1 15
Give a tablet, repeated in thirty to sixty minutes, as an emetic in croup, with draughts of warm water.		
IPECAC POWD.	5 grs.	3 00
Emetic.		
One tablet with warm water, as an emetic; if it fails to act promptly repeat in an hour, with the addition of a teaspoonful of mustard.		
IPECAC, FL. EXT.	1 min.	1 50
One every two hours, for colds and to promote sweating in fevers; one after meals as a hepatic stimulant; useful in bilious headache.		
IPECAC WINE.	1 min.	75
One after meals as a stomachic stimulant; one every half hour for headache of rheumatic patients.		
IPECAC WINE.	2 min.	75
One three times a day for gastric and duodenal catarrh, and irritative diarrhœa, in children.		
IPECAC WINE.	5 min.	1 00
One, repeated in two hours, for sick-headache and migraine, and as a stimulant to the alimentary canal.		
IRON ALBUMINATE.	3 grs.	2 25
Nutritive tonic.		
One after meals, for anæmia in children.		
IRON ALBUMINATE.	5 grs.	3 50
Three or four daily for anæmia, chlorosis, and after wasting disease.		
IRON ALBUMINATE COMP.	Iron Albuminate.... 5 gr. / Modified nutritive tonic. Acid Arsenous... 1-40 gr. }	3 50
One after each meal, as a tonic.		
IRON & ALOES COMP.	{ Strych. Sulph.... 1-60 gr. } Laxative tonic. { Acid Arsenous... 1-30 gr. } { Ext. Aloes..... 1-12 gr. } { Iron Powd..... 1 gr. }	1 10
One or two at night, or every second night, for constipation with intestinal atony, in anæmia.		
IRON ARSENIATE.	1-25 gr.	75
Tonic.		
One, three times daily for tonic.		
IRON ARSENIATE.	1-16 gr.	75
Three to four a day, in simple anæmia and chlorosis.		
IRON ARSENIATE.	1-10 gr.	75
One after each meal, as a hæmal tonic.		

IRON, ARSENIC & BRUCIA (Dr. Clymer.) Tonic and hæmætic.	{ Iron Powder.....1-10 gr. } { Acid Arsen.....1-100 gr. } { Brucia.....1-100 gr. }	75
Give one about an hour after eating, as a stomachic and general tonic.		
IRON, ARSENIC AND BRUCIA No. 2.....	{ Iron Powder.....1-10 gr. } { Acid Arsen.....1-100 gr. } { Brucia.....1-33 gr. }	75
Give one about an hour after eating, as a stomachic and general tonic.		
IRON, ARSENIC AND BRUCIA No. 3.....	{ Iron Powder.....1 gr. } { Acid Arsen.....1-100 gr. } { Brucia.....1-100 gr. }	75
Give one about an hour after eating, as a stomachic and general tonic.		
IRON & ARSENIC COMP. (Dr. Clymer.)	{ Iron Powder.....1 gr. } { Acid Arsen.....1-100 gr. } { Ignatia Powd....1-40 gr. }	75
Give one about an hour after eating, as a stomachic and general tonic.		
IRON & ARSENIC COMP. No. 2 (Dr. Clymer.)	{ Iron Powder.....1 gr. } { Acid Arsenous...1-50 gr. } { Ignatia Powd....1-40 gr. }	75
Give one about an hour after eating, as a stomachic and general tonic.		
IRON, ARSENIC AND NUX ...	{ Iron Powder.....1 gr. } { Acid Arsen.....1-100 gr. } { Ext. Nux Vomica 1-8 gr. }	75
A tablet after meals, increasing until three are taken at a dose, for anæmia with palpitation, dyspepsia, and reflex nervous symptoms.		
IRON, ARSENIC AND STRYCH- NINE Iron tonic.	{ Iron Powder.....1 gr. } { Acid Arsen.....1-100 gr. } { Strych. Sulph....1-60 gr. }	75
One tablet three times a day, ten minutes before meals; a useful tonic in all kinds of anæmia, with loss of appetite.		
IRON, ARSENIC AND STRYCH- NINE No. 2	{ Iron Powder.....1 gr. } { Acid Arsen.....1-50 gr. } { Strych. Sulph....1-60 gr. }	75
One after each meal and at bed hour, as a general tonic and for loss of appetite.		
IRON, ARSENIC AND STRYCH- NINE No. 3	{ Iron Powder.....1 gr. } { Acid Arsen.....1-50 gr. } { Strych. Sulph....1-100 gr. }	75
One after each meal and at bed hour, for general tonic, and loss of appetite.		
IRON, ARSENIC AND STRYCH- NINE No. 4	{ Iron Powd.....2 gr. } { Acid Arsenous...1-80 gr. } { Strych. Sulph....1-60 gr. }	1 10
One after each meal and at bed hour, for general tonic, loss of appetite and anæmia.		
IRON, ARSENIC AND STRYCH- NINE No. 5	{ Iron Powder.....1 gr. } { Acid Arsenous...1-20 gr. } { Strych. Sulph....1-30 gr. }	75
One after each meal and at bed hour, for general tonic, loss of appetite and anæmia.		
IRON CARBON. FRASER'S No. 1..... Tonic.	{ Ferri Carb.....2 gr. } { Sacch. Alb.....1 gr. }	1 15
One after meals, for anæmic and convalescent children.		
IRON CARBON. FRASER'S No. 2.....	{ Ferri Carb.....5 gr. } { Sacch. Alb.....2 1-2 gr. }	1 50
One every three hours, for tonic and after wasting diseases		

When ordering or prescribing

IRON POWDER	1-50 gr.....	75
One every three hours, increasing every day by one till three or four are taken at a dose.		
IRON POWDER	1-10 gr.....	75
One every three hours for anæmia in childhood.		
IRON POWDER	1-5 gr.....	75
One after meals, for pale and sickly children.		
IRON POWDER	1-2 gr.....	75
One, three or four times a day, for anæmia, amenorrhœa and allied disorders.		
IRON POWDER	1 gr.....	75
A tablet three or four times a day, for anæmia, anæmic headache and in convalescence.		
IRON POWDER	2 grs.....	1 10
Three to four daily, for anæmia and various cachexiæ.		
IRON PYROPHOSPHATE	1-10 gr.....	75
One every two hours for anæmic diseases of childhood, especially rhachitis.		
IRON PYROPHOSPHATE	1-5 gr.....	75
One to three after meals, for malnutrition of childhood.		
IRON PYROPHOSPHATE	1-2 gr.....	75
One or two after meals, for rhachitic and tubercular disease in childhood.		
IRON PYROPHOSPHATE	1 gr.....	1 10
A tablet three times a day, for a tonic in malnutrition and catamenial disorders.		
IRON PYROPHOS COMP.	<div> <div>Iron Pyrophos.....1 gr.</div> <div>Quinine Sulph.....1-2 gr.</div> <div>Strych. Sulph.....1-60 gr.</div> <div>Acid Arsenous.....1-30 gr.</div> </div>	1 00
Modified iron tonic.		
One after meals, as a hæmatic and alterative tonic.		
IRON, QUININE & ALOES CO.	<div> <div>Iron Powd.....2 gr.</div> <div>Quinine Sulph.....2 gr.</div> <div>Strych. Sulph.....1-40 gr.</div> <div>Acid Arsenous.....1-40 gr.</div> <div>Aloes Powd.....1-8 gr.</div> <div>Per 100.....</div> </div>	4 00
(Dr. Duncan.)		
One every three hours in chronic malaria.		
IRON, QUININE AND STRYCH-NINE	<div> <div>Iron Powder.....1-2 gr.</div> <div>Quinine Sulph.....1-2 gr.</div> <div>Strych. Sulph.....1-120 gr.</div> </div>	1 50
Tonic, antiperiodic.		
One or two after meals, as an antimalarial and general tonic.		
IRON & STRYCHNINE	<div> <div>Iron Powder.....1 gr.</div> <div>Strych. Sulph.....1-60 gr.</div> </div>	75
Tonic.		
Two or three tablets a day, in nervous and anæmic diseases of adolescence.		
IRON & STRYCHNINE COMP	<div> <div>Iron Powder.....1-2 gr.</div> <div>Quinine Sulph.....1-2 gr.</div> <div>Acid Arsen.....1-100 gr.</div> <div>Strych. Sulph.....1-120 gr.</div> </div>	1 50
(Dr. Morrill.)		
Tonic, antiperiodic.		
One or two, according to age, three times a day, after meals. Used in chronic malarial poisoning, malarial anæmia, neuralgias, chlorosis and the amenorrhœa of anæmia.		
IRON & STRYCHNINE CIT	1-2 gr.....	90
General nerve tonic.		
One after meals, as a hæmatic and nerve tonic in chorea, epilepsy, cerebral and spinal anæmia, and hysteria.		

IRON & STRYCHNINE CIT.	1 gr.	1 15
One after meals, as a hæmatic and nerve tonic in chorea, epilepsy cerebral and spinal anæmia, and hysteria.		
JALAP	1-2 gr.	75
One every half hour for five doses, then every hour until laxative effect is produced.		
JALAP	1 gr.	1 10
One every half hour for five doses, then every hour until laxative effect is produced.		
KALI COMP.	<div> <div>Kali Bichrom.1-100 gr. }</div> <div>Mercury Biniod. 1-100 gr. }</div> </div>	75
One every two hours, for sore throat from glandular irritation.		
KINO COMP.	<div> <div>Kino Powd.3 3-4 gr. }</div> <div>Cinnamon Powd.1 gr. }</div> <div>Opium Powd.1-4 gr. }</div> </div>	2 50
One every three or four hours, as astringent, in diarrhœa.		
KRAMERIA COMP.	<div> <div>Fl. Ext. Krameria . . .1 min. }</div> <div>Bismuth Subnit.1 gr. }</div> <div>Opium Powd.1-4 gr. }</div> <div>Licorice R't Po.1-4 gr. }</div> </div>	1 00
(Dr. Applegate.) Diarrhœa.		
One every hour, for diarrhœa and dysentery; decrease to one every three or four hours as symptoms ameliorate; valuable also in intestinal inflammation, used in conjunction with, or following salines.		
KRAMERIA COMP. No. 2.	<div> <div>Fl. Ext. Krameria 1-2 min. }</div> <div>Bismuth Subnit.1 gr. }</div> <div>Pulv. Opii1-25 gr. }</div> </div>	1 00
One tablet every two or three hours powdered in sugar, for the summer diarrhœa of children; allow only boiled milk until the movements become normal.		
LACTO FERRIC	<div> <div>Iron Lactate.1 gr. }</div> <div>Cinchonia Sulph.1 gr. }</div> <div>Ext. Hyoseyam.1-5 gr. }</div> <div>Acid Arsenous.1-30 gr. }</div> <div>Strych. Sulph.1-60 gr. }</div> </div>	1 50
Tonic.		
A tablet three or four times a day, as a tonic in malnutrition and anæmia with neuroses and ataxia.		
LARYNGITIS	<div> <div>Potass. Brom.1 1-4 gr. }</div> <div>Potass. Chlorate.5-8 gr. }</div> <div>Ipecac Powd.1-12 gr. }</div> <div>Ext. Licorice1 1-4 gr. }</div> <div>Squills Powd.1-2 gr. }</div> </div>	1 15
One every hour, for continued cough, with tickling in the throat.		
LAXATIVE	<div> <div>Aloin1-4 gr. }</div> <div>Ext. Belladonna.1-8 gr. }</div> <div>Strych. Sulph.1-60 gr. }</div> <div>Ipecac Pulv.1-16 gr. }</div> </div>	1 15
One or two at night, as a stimulant to the alimentary canal and to increase peristalsis.		
LAXATIVE No. 2.	<div> <div>Quinine Sulph.1 gr. }</div> <div>Ext. Rhubarb.1 gr. }</div> <div>Ipecac Powd.1-8 gr. }</div> <div>Capsicum Powd.1-4 gr. }</div> </div>	3 00
(Dr. Hauck.)		
One or two at a dose every two or three hours, as required for catarrh of the stomach, and dyspepsia, combined with mild forms of constipation.		

Please specify Fraser's Tablets

LAXATIVE No. 3 (Dr. Burleigh.)	{ Aloes.....1 gr. }	2 00
	{ Ext. Belladonna.3-20 gr. }	
	{ Ext. Physostig.3-20 gr. }	
	{ Ext. Nux Vomica.1-5 gr. }	
	{ Ext. Cascara.....1-5 gr. }	

One or two tablets at night, repeated the next morning if necessary, for constipation.

LAXATIVE No. 4 (Dr. A. M. Thomas.)	{ Ext.Cascara Sag.1-2 gr. }	2 50
	{ Sodium Bicarb.1-4 gr. }	
	{ Hydrastin.....1-200 gr. }	
	{ Ext. Nux Vom.1-8 gr. }	
	{ Ext. Belladonna.1-16 gr. }	
	{ Oil Peppermint.1-40 min. }	

One to four at night, as required as a laxative. Can be taken indefinitely without producing habit.

LEAD ACETATE1 gr.	75
-------------------------------------	----

For solution.

One tablet dissolved in a tablespoonful of water makes a solution of about 1-250; for external use and as an injection.

LEAD & BISMUTH COMP	{ Lead Acetate.....1 gr. }	2 25
	{ Bismuth Subnit.....2 gr. }	
	{ Camphor.....1-8 gr. }	
	{ Opium Powd.....1-4 gr. }	

Give a tablet every hour for three or four doses, in bilious and flatulent diarrhoea; one after meals for gastric catarrh.

LIME & IRON PHOSPHATES (Dr. Hubbard.)	{ Lime Phosphate....2 gr. }	1 50
	{ Iron Phosphate....2 gr. }	
	{ Sugar.....1 gr. }	
	{ Oil Wintergr'n.1-10 min. }	

One tablet to a child; two to an adult, after each meal, as a reconstructive tonic.

LIME WATER	1 00
-------------------------	------

For the preparation of Lime Water, add six tablets to a half pint of boiled water in a clean bottle, and allow to stand with occasional agitation until the tablets disintegrate. Shake again and set aside until the excess of lime has settled and the mixture becomes clear. Decant the clear liquid if desired and keep well bottled for use.

LITHIUM BENZOATE1 gr.	1 75
---	------

One every three hours, for secondary rheumatic affections.

LITHIUM CARBONATE1 gr.	1 70
--	------

Antilithic.

One every hour, for acute rheumatism; one to three after meals to hasten the excretion of uric acid.

LITHIUM CARBONATE5 grs.....	3 50
--	------

Anthilitic.

One or two after each meal, for gout, arthritis deformans, gravel and vesical calculus.

LITHIUM CARBONATE (EFFERVESCENT.)	3 50
---	------

Each tablet represents three grains lithium salt.

A tablet three or four times a day, in a glass of water, for rheumatism and gout and the uric acid diathesis.

LITHIUM SALICYLATE5 grs.....	6 00
---	------

Antiseptic alkaline.

One, three or four times a day, for the uric acid diathesis, muscular rheumatism and dyspepsia of gouty subjects.

When ordering or prescribing

LUPULIN & BROMIDE CO....	Lupulin.....	1-2 gr.	} 3 25
	Scutellarin.....	1-2 gr.	
	Ergotin.....	1-4 gr.	
	Atropine Sulph.....	1-2000 gr.	
	Zinc Bromide.....	1-10 gr.	

One or two ; three or four times a day, as a general nerve sedative.

MALARIA (Dr. Clarke.)	Quinine Sulph.....	3 gr.	} 5 50
	Capsicum Powd.....	1 gr.	
	Opium Powd.....	1-4 gr.	

Two to four tablets at nine in the forenoon, one at three P. M. and one at bedtime ; for remittent and malarial fever.

MANGANESE BINOXIDE.....	1-10 gr.	75
-------------------------	----------	----

One tablet, crushed in a teaspoonful of sweetened water, three or four times a day, for an infant with anæmia.

MANGANESE BINOXIDE.....	1 gr.	75
-------------------------	-------	----

One after meals, for reconstructive tonic, where iron does not agree.

MANGANESE BINOXIDE.....	2 grs.	1 25
-------------------------	--------	------

One tablet every three hours, for a general tonic, and as a mild emmenagogue.

MANGANESE BINOXIDE.....	3 gr.	1 00
-------------------------	-------	------

One every three hours, beginning a few days before the period, for suppressed or scanty menstruation.

MANGANESE BINOXIDE.....	5 gr.	2 00
-------------------------	-------	------

One, three times a day, as an emmenagogue and regulator ; has also been used with good results in amenorrhœa.

MENORRHAGIC. (Dr. Hirst.)	Acid Gallic.....	3 gr.	} 5 00
	Ergotin.....	1 gr.	
	Hydrastin.....	1-2 gr.	

One or two, every two or three hours as required.

MERCURIUS COMP Throat troubles.	Merc. Iod. Red.....	1-100 gr.	} 75
	Belladonna Po.....	1-100 gr.	
	Kali Bichrom.....	1-100 gr.	

One every hour or two, for red and inflamed throat.

MERCURIAL TONIC.....	Mercury Protiod.....	1-8 gr.	} 1 10
	Iron Citrate.....	1-2 gr.	

One after meals, as a hæmatic and alterative tonic for chronic diseases of the blood and skin.

MERCURY, BELLADONNA CO Useful in throat troubles.	Merc. Iod. Red.....	1-200 gr.	} 75
	Ipecac Powd.....	1-20 gr.	
	Belladonna Po.....	1-100 gr.	

A tablet every hour or two, for sore throat with swollen glands.

MERCURY & CHARCOAL..... (Dr. R. Tilley.)	Mercury Protiod.....	1-4 gr.	} 75
	Carbo Veg.....	1-10 gr.	

Alterative.

One, half an hour after meals, for chronic skin and visceral disease. The Carbo Veg is added to change the color.

MERCURY & CHARCOAL No. 2	Mercury Protiod.....	1-6 gr.	} 75
	Carbo Veg.....	1-10 gr.	

One after meals, as a general tonic and alterative.

MERCURY & HYOSCYAMUS.....	Mercury Protiod.....	1-4 gr.	} 1 10
	Ext. Hyoscyam.....	1-4 gr.	

One every three hours, as a general alterative.

MERCURY IODIDE COMP..... (Dr. Curtis.)	Merc. Iod. Red.....	1-32 gr.	} 75
	Ipecac Powd.....	1-8 gr.	

One or two after meals, for the continued treatment of syphilis.

MERCURY & IRON	Mercury Protiod. 1-4 gr. {	1 10
Ferruginous alterative.	Iron Powder.....1 gr. }	
A tablet with meals, as an alterative and hæmatic tonic, for chronic diseases and anæmia.		
MERCURY & OPIUM	Mercury Protiod. 1-5 gr. {	95
	Opium Powd.....1-12 gr. }	
One three times a day, for tonic syphilitic treatment.		
MERCURY & OPIUM CO.	Merc. Protiod.1-4 gr. {	1 00
	Opium Powd.....1-12 gr. }	
	Charcoal.....1-10 gr. }	
One, three times a day, for tonic syphilitic treatment.		
MERCURY & POTASS. IODIDE	Merc. Iod. Red.....1 gr. {	2 25
(For antiseptic use.)	Potass. Iodide.....1 gr. }	
One tablet dissolved in four tablespoonfuls of water makes a solution of about 1 to 1000.		
MERCURY & RHUBARB	Blue Mass.....2 gr. {	1 50
Alterative.	Comp. Rhubarb Pil. 1 gr. }	
Give one after each meal, as an alterative.		
MERCURY PROTIODIDE	1-1000 gr.	75
One every hour, for the vomiting and diarrhœa of infantile gastro-enteritis.		
MERCURY PROTIODIDE	1-500 gr.	75
Give one tablet every three hours, to an infant, to promote intestinal antiseptis.		
MERCURY PROTIODIDE	1-200 gr.	75
One every two hours, to an infant, for hepatic stimulant and aid to digestion.		
MERCURY PROTIODIDE	1-100 gr.	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
MERCURY PROTIODIDE	1-50 gr.	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
MERCURY PROTIODIDE	1-25 gr.	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
MERCURY PROTIODIDE	1-20 gr.	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
MERCURY PROTIODIDE	1-16 gr.	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
MERCURY PROTIODIDE	1-10 gr.	75
One every three hours, as a hepatic stimulant; also used in continued constitutional treatment.		
MERCURY PROTIODIDE	1-8 gr.	75
One or two every three hours, in the tonic treatment of syphilis.		
MERCURY PROTIODIDE	1-6 gr.	75
One or two every three hours, in the tonic treatment of syphilis.		
MERCURY PROTIODIDE	1-5 gr.	75
One or two every three hours, in the tonic treatment of syphilis.		
MERCURY PROTIODIDE	1-4 gr.	75
One every three hours, to produce rapid mercurialization; decrease the dose as soon as possible.		

MERCURY PROTIODIDE1-3 gr.....	95
One every three hours, to produce rapid mercurialization; decrease the dose as soon as possible.	
MERCURY PROTIODIDE1-2 gr.....	1 20
A tablet every three hours, with large doses of potass. iodide, to control cerebral symptoms in syphilis; decrease the mercury as symptoms improve.	
MERCURY PROTIODIDE1 gr.....	1 85
A tablet every three hours, with large doses of potass. iodide, to control cerebral symptoms in syphilis; decrease the mercury as symptoms improve.	
MERCURY SALICYLATE1-4 gr.....	1 40
One every hour, for a rapid effect without severe irritation; lessen the dose as the result is obtained.	
MERCURY SALICYLATE1-3 gr.....	1 60
One every two hours, for a rapid effect without severe irritation; lessen the dose as the result is obtained.	
MERCURY TANNATE1-2 gr.....	1 50
One every two hours, for a rapid effect without severe irritation; lessen the dose as the result is obtained.	
MERCURY WITH CHALK1-100 gr.....	75
One, in a little water on the end of a spoon, repeated every hour for infantile cholic with profuse diarrhœa.	
MERCURY WITH CHALK1-40 gr.....	75
One every three hours to a child, for colic with profuse diarrhœa.	
MERCURY WITH CHALK1-20 gr.....	75
One every hour for an adult, for hepatic corrective, until ten or fifteen have been taken.	
MERCURY WITH CHALK1-10 gr.....	75
One every hour, for hepatic corrective, until effect is produced.	
MERCURY WITH CHALK1-3 gr.....	75
One every two hours, for tonic to liver, and corrective in bilious diarrhœa until effect is produced; as an alterative, one three times a day.	
MERCURY WITH CHALK1-2 gr.....	75
One every two hours, for tonic to liver, and corrective in bilious diarrhœa—as an alterative, one three times a day.	
MERCURY WITH CHALK1 gr.....	75
One tablet repeated every three hours until effect is produced, as hepatic alterative with laxative.	
MERCURY YELLOW OXIDE1-1000 gr.....	75
One every hour, for a child with indigestion and diarrhœa, to control septic conditions of the intestine.	
MERCURY YELLOW OXIDE1-200 gr.....	75
One tablet every two or three hours, for dysentery; one after meals as an alterative in chronic disease.	
MERCURY YELLOW OXIDE1-100 gr.....	75
A tablet after meals, for chronic diseases of the skin and lymphatics; one every hour to secure immediate constitutional effects.	
MIGRAINE	2 00
<div style="display: flex; align-items: center;"> <div style="flex: 1;"> <div style="font-size: 2em; margin-right: 10px;">{</div> <div> Acetanilid.....2 gr. Camph. Monobr.....1-2 gr. Caffeine Citrate.....1-2 gr. </div> </div> </div>	
Two tablets, repeated every hour, for four or five doses, for neuralgic and nervous headache.	
MIGRAINE No. 2	3 00
<div style="display: flex; align-items: center;"> <div style="flex: 1;"> <div style="font-size: 2em; margin-right: 10px;">{</div> <div> Acetanilid.....3 gr. Camph. Monobrom.....2 gr. Caffeine Cit.....1 gr. </div> </div> </div>	
One, repeated every hour, for three doses, then every two hours; as a relief to migraine, supraorbital and frontal.	

MIGRAINE No. 3	{ Acetanilid3 1-2 gr. } { Caffeine Alk1-2 gr. } { Sodium Bicarb.....1 gr. }	2 00
-----------------------------	---	------

One, repeated in an hour, for sick headache, and pains in the head with fever, may be continued for three doses, then stop and begin again in three hours if not relieved.

MILK	{ Sugar of Milk.....26 gr. } { Calc. Lactophos...1-6 gr. } { Calc. Carbonate...1-2 gr. } { Sodium Bicarb....1-2 gr. } { Potass. Bicarb....1-12 gr. } { Sodium Chloride...1-6 gr. }	2 25
-------------------	---	------

Four tablets to a bottle of milk, as a reconstructive tonic for bottle fed babies.

MIXED TREATMENT	{ Potass. Iodide,5 gr. } { Corrosive Sub....1-30 gr. } { Syr. Sars. Co30 min. }	5 65
------------------------------	---	------

Alterative and antisyphilitic. Begin with three a day, and increase according to the necessities of the case.

MORPHINE & ACONITE	{ Morph. Sulph....1-32 gr. } { Ext. Aconite....1-100 gr. }	1 10
-------------------------------------	---	------

MORPHINE & ACONITE No. 2	{ Morph. Sulph....1-8 gr. } { Ext. Aconite....1-25 gr. }	2 00
---	---	------

MORPHINE & BELLADONNA	{ Morph. Sulph....1-6 gr. } { Ext. Belladonna...1-8 gr. }	2 40
--	--	------

MORPHINE COMP. No. 1	{ Morph. Sulph....1-8 gr. } { Atropine Sulph.1-150 gr. }	2 00
-----------------------------------	---	------

MORPHINE COMP. No. 2	{ Morph. Sulph....1-4 gr. } { Atropine Sulph.1-120 gr. }	3 35
-----------------------------------	---	------

MORPHINE COMP. No. 3	{ Morph. Sulph....1-8 gr. } { Atropine Sulph.1-200 gr. }	2 00
-----------------------------------	---	------

MORPHINE COMP. No. 4	{ Morph. Sulph....1-4 gr. } { Atropine Sulph.1-150 gr. }	3 35
-----------------------------------	---	------

MORPHINE SULPHATE	1-100 gr.....	75
--------------------------------	---------------	----

MORPHINE SULPHATE	1-50 gr.....	95
--------------------------------	--------------	----

MORPHINE SULPHATE	1-20 gr.....	1 00
--------------------------------	--------------	------

MORPHINE SULPHATE	1-10 gr.....	1 30
--------------------------------	--------------	------

MORPHINE SULPHATE	1-10 gr.....	1 50
--------------------------------	--------------	------

MORPHINE SULPHATE	1-8 gr.....	1 85
--------------------------------	-------------	------

MORPHINE SULPHATE	1-6 gr.....	2 25
--------------------------------	-------------	------

MORPHINE SULPHATE	1-4 gr.....	3 20
--------------------------------	-------------	------

MORPHINE SULPHATE	1-3 gr.....	3 75
--------------------------------	-------------	------

MORPHINE SULPHATE	1-2 gr.....	5 00
--------------------------------	-------------	------

MUSCARIN SULPHATE	1-100 gr.....	7 00
--------------------------------	---------------	------

A tablet every three or four hours stimulates the cutaneous, salivary and intestinal glands, and the intra cardiac inhibitory centres; in scar latina, rheumatism and respiratory diseases.

NAPHTALIN, Pure	5 grs.....	2 00
------------------------------	------------	------

Antiseptic.

A tablet three to six times a day for the diarrhoea and elevated temperature in typhoid; one to three as an anthelmintic, followed by a dose of castor oil.

NAUSEA No. 2	{ Bismuth Subnit.....2 gr. } { Cerium Oxalate...2 gr. } { Cocaine Hydro...1-12 gr. }	4 75
---------------------------	--	------

One every hour until stomach is settled, or ten doses are taken. Useful for car sickness or sea sickness.

NAUSEA No. 3 } Cerium Oxalate....5 gr. (2 25
 (Dr. Guernsey.) } Ipecac Powd....1-10 gr. }

One every three or four hours, for vomiting of pregnancy.

NERVINE..... } Camphor.....1 gr. } 1 50
 } Ext. Hyoscyam...1-2 gr. }
 } Valerian Powd...1-2 gr. }

One every half hour for three doses, then every two hours until nervousness is relieved.

NERVINE No. 2..... } Lupulin.....1 gr. } 2 00
 (Dr. Lee.) } Zinc Phosphide..1-16 gr. }
 } Sodium Bromide..2 1-2 gr. }
 } Gelsemin1-20 gr. }
 } Acid Arsenous...1-20 gr. }

Two tablets, three times a day, for irritability of the nervous system and neurasthenia.

NERVE TONIC No. 1..... } Zinc Phosphide..1-10 gr. } 1 40
 (Dr. Westbrook.) } Ext. Nux Vom....1-4 gr. }
 } Iron Powder.....1 gr. }

One, three times a day, may be given as long as improvement continues or until stomach is nauseated; if nausea ensues, may be discontinued for one week, then repeated.

NERVE TONIC No. 2..... } Zinc Phosphide..1-10 gr. } 1 30
 (Dr. Westbrook.) } Ext. Nux Vom....1-4 gr. }
 } Acid Arsenous...1-20 gr. }

One, three times a day, may be given as long as improvement continues. If nausea or redness of the eyelids, or pains or fulness of the stomach occur, discontinue the tablet for a week, then continue.

NERVE TONIC No. 3..... } Zinc Phosphide..1-10 gr. } 1 75
 (Dr. Westbrook.) } Ext. Nux Vom....1-4 gr. }
 } Iron Powd.....2 gr. }

One, three times a day, may be given as long as improvement continues or until stomach is nauseated; if nausea ensues, may be discontinued for one week, then repeated.

NERVE TONIC No. 4..... } Zinc Phosphide..1-10 gr. } 1 75
 (Dr. Westbrook.) } Ext. Nux Vom....1-4 gr. }
 } Iron Powder.....2 gr. }
 } Acid Arsenous...1-20 gr. }

One, three times a day, may be given as long as improvement continues. If nausea or redness of the eyelids, or pains or fulness of the stomach occur, discontinue the tablet for a week, then continue.

NERVE TONIC No. 5..... } Zinc Phosphide...1-8 gr. } 1 30
 } Ext. Nux Vom....1-8 gr. }
 } Ext. Can. Indica..1-8 gr. }

One with meals, for nervous affections with exalted reflexes and excitability.

NERVE TONIC No. 6..... } Ext. Cannab. Ind..1-8 gr. } 2 25
 } Hyoscyamia....1-400 gr. }
 } Zinc Phosphide..1-10 gr. }

Three or four tablets a day, for chorea, neurasthenia and hysteria.

NERVE TONIC No. 7..... } Hyoscyamine...1-400 gr. } 2 50
 } Zinc Phos.....1-10 gr. }
 } Strych. Nit.....1-20 gr. }

One, three times a day, for disorders of the nerves dependent upon anæmia and innutrition of the cord.

NEURALGIC..... } Quinine Sulph....1-2 gr. } 1 70
 (Dr. Dunlap.) } Morph. Sulph....1-80 gr. }
 } Strych. Sulph....1-120 gr. }
 } Acid Arsenous...1-80 gr. }
 } Ext. Aconite.....1-8 gr. }

One every hour until eight are taken, if necessary. For neuralgia and neuralgic headache.

NEURALGIC No. 2 (Dr. Henry J. Kenyon.)	{ Zinc Phosphide...1-16 gr.	1 50
	{ Strych. Sulph....1-60 gr.	
	{ Ext. Can. Indica...1-8 gr.	
	{ Sodium Arsen....1-20 gr.	
	{ Aconitine.....1-400 gr.	

NEURALGIC No. 3 (Dr. Miller.)	{ Acetanilid.....2 1-2 gr.	2 35
	{ Camph.Monobr. 2 1-2 gr.	

A tablet every two hours for three doses, as a sedative and anodyne in general and localized neuralgias.

NEURALGIC No. 4 (Dr. Gustavus Elliot.)	{ Fl. Ext. Bellad...1-6 min.	1 50
	{ Fl. Ext. Aconite.1-2 min.	
	{ Fl.Ext.Gelsem.1 1-3 min.	

One every hour or every second hour, for painful affections of the nerves, not due to an organic lesion.

NEURALGIC HEADACHE ... (Dr. Myers.)	{ Sodium Bromide....5 gr.	2 10
	{ Caffeine Citrate...1-2 gr.	
	{ Antifebrin.....1 gr.	
	{ Ext. Hyoseyam...1-2 gr.	
	{ Morph.Sulph....1-50 gr.	

A tablet, repeated in half an hour, for migraine, headache and facial neuralgia.

NEURASTHENIA (Dr. Krauss.)	{ Lithium Carb.....3 gr.	3 45
	{ Iron Phosphate...1-3 gr.	
	{ Ext. Nux Vomica...1 gr.	
	{ Sodium Arsen....1-27 gr.	

One to three tablets after meals, for neurasthenia and lithæmia with predominant nervous symptoms.

NEUROTIC (Dr.North.)	{ Tr. Benzoin Co....2 min.	2 45
	{ Tr. Lavender Co...4 min.	
	{ Tr. Valer. Amm...4 min.	

Give one tablet every ten minutes until relieved, for hysteria and nervousness of women.

NEUTRALIZING	{ Pulv. Rhei.....1 gr.	1 50
	{ Sodium Bicarb....3 gr.	
	{ Magnesium Calc...1 gr.	
	{ Pulv. Cardamom...1-8 gr.	
	{ Oil Menth. Pip...1-8 gr.	

One or two tablets half an hour after eating, for gastralgia and flatulence, and sour eructations.

NITROGLYCERIN1-200 gr.....	75
---	----

Heart tonic and in asthma.

One when face is pale, for fainting, neuralgia of heart or heart failure. Never give if face is flushed.

NITROGLYCERIN1-150 gr.....	75
---	----

A tablet, repeated in an hour if necessary, for pallor, headache and syncope; one three times a day for epilepsy, catalepsy and angina-pectoris.

NITROGLYCERIN1-100 gr.....	75
---	----

One when face is pale, for fainting, neuralgia of heart or heart failure. Never give if face is flushed.

NITROGLYCERIN1-50 gr.....	75
--	----

One tablet, repeated in two or three hours if necessary, for cardiac syncope, epileptiform seizure, extreme asthmatic paroxysms, angina pectoris, and parturient eclampsia.

NITROGLYCERIN CO. No. 1 ...	{ Nitroglycerin...1-100 gr.	1 25
	{ Tr. Digitalis.....2 min.	

One, three times a day for feeble heart and cerebral anæmia; for epilepsy, increase by one tablet every three weeks.

NITROGLYCERIN CO. No. 2	{ Nitroglycerin. . . 1-100 gr. } { Tr. Strophanthus. . 2 min. }	1 50
A tablet three times a day ; for nephritis with pulse of great tension, cardiac weakness, and the anæmic form of migraine.		
NUX VOMICA EXT.	1-100 gr.	75
One every two hours, as a tonic stomachic in dyspepsia.		
NUX VOMICA EXT.	1-50 gr.	75
One every two hours, as a tonic stomachic in dyspepsia.		
NUX VOMICA EXT.	1-10 gr.	75
One after each meal, as a general nerve tonic and a tonic to stomach in dyspepsia.		
NUX VOMICA EXT.	1-8 gr.	75
One after each meal, as a general nerve tonic and a tonic to stomach in dyspepsia.		
NUX VOMICA EXT.	1-4 gr.	75
One, three times a day, as a general nerve tonic and to promote intestinal peristalsis.		
NUX VOMICA FL. EXT.	1 min.	1 10
One after meals, for tonic to stomach in dyspepsia.		
NUX VOMICA POWD.	1-1000 gr.	75
One three times a day for a child, as a tonic and stimulant.		
NUX VOMICA POWD.	1-500 gr.	75
Give one or two every two hours, to increase the flow of gastric juice and aid peristalsis.		
NUX VOMICA POWD.	1-200 gr.	75
Give one every two hours, as a general tonic and an aid to digestion.		
NUX VOMICA POWD.	1-100 gr.	75
Give one every two hours, as a general tonic and an aid to digestion.		
NUX VOMICA POWD.	1-50 gr.	75
Give one every two hours, as a general tonic and an aid to digestion.		
NUX VOMICA POWD.	1-20 gr.	75
A tablet every two hours, for dyspepsia and atonic diarrhœa of hot weather.		
NUX VOMICA POWD.	1-10 gr.	75
One after each meal, repeated in an hour if necessary, as an aid to digestion and tonic to the stomach.		
NUX VOMICA POWD.	1-8 gr.	75
One after each meal, repeated in an hour if necessary, as an aid to digestion and tonic to the stomach.		
NUX VOMICA POWD.	1-4 gr.	75
One after each meal, as a nerve tonic and aid to digestion.		
NUX VOMICA POWD.	1-2 gr.	75
One after meals, for atonic dyspepsia, gastric or intestinal.		
NUX VOMICA COMP.	{ Ext. Nux Vom. . . 1-64 gr. } { Ext. Belladonna . . 1-32 gr. } { Ipecac Powd . . . 1-10 gr. } { Aloin 1-8 gr. } { Podophyllin. . . . 1-4 gr. }	1 50
Tonic cathartic.		
One to four at night, for constipation depending upon loss of function of the liver and intestines.		
NUX VOMICA COMP. No. 2.	{ Ext. Nux Vom. . . 1-16 gr. } { Acid Arsenous. . . 1-80 gr. } { Pepsin Pure. . . . 1-32 gr. } { Aromatics. q. s. }	75
One, three or four times a day, for anorexia, dyspepsia and constipation.		

NUX VOMICA COMP. No. 3....	{ Ext. Nux Vom....1-20 gr. }	75
	{ Sulphur.....1-20 gr. }	
	{ Acid Arsenous...1-100 gr. }	

One or two after meals, for chronic skin diseases dependent upon dyspepsia.

NUX & ARSENIC.....	{ Nux Vom. Po....1-10 gr. }	75
	{ Acid Arsenous...1-50 gr. }	

Tonic, antiperiodic.

One after meals, as a tonic in malarial indigestion.

NUX & BRYONIA.....	{ Nux Vom. Po....1-100 gr. }	75
	{ Bryonia Powd...1-100 gr. }	

For a child, give one every two hours, as a tonic to the stomach and an aid to digestion, with slight laxative action.

NUX & BRYONIA No. 2.....	{ Tr. Nux Vom.....1 min. }	1 10
	{ Tr. Bryonia.....1 min. }	

One every two or three hours, as a tonic to the stomach and aid to digestion.

NUX & CANTHARIDES.....	{ Nux Vom. Po....1-100 gr. }	75
	{ Cantharides....1-100 gr. }	

A tablet three times a day, for impotence and spermatorrhœa; best combined with iron.

NUX & CARBO VEG.....	{ Nux Vom. Po....1-100 gr. }	75
	{ Carbo Veg.....1-10 gr. }	

Tonic, absorbent.

One after meals, for atonic gastric disorders in children, and those due to fermentation.

NUX & CARBO VEG. No. 2.....	{ Nux Vom. Po....1-2 gr. }	75
	{ Carbo Veg.....1-4 gr. }	

A tablet three times a day, for chronic catarrhal inflammation of the stomach, with flatulence.

NUX & COLOCYNTH CO.....	{ Ext. Belladonna...1-8 gr. }	2 85
	{ Ext. Nux Vom....1-4 gr. }	
	{ Ext. Colocy. Co. 1 1-2 gr. }	
	{ Sodium Bicarb.....1 gr. }	

One every night, for habitual constipation and disordered hepatic function.

NUX & PEPSIN.....	{ Nux Vom. Po....1-100 gr. }	75
	{ Pepsin Pure.....1-10 gr. }	

(Dr. Skiff.)

Tonic, antidyspeptic.

Two tablets after each meal, for simple indigestion. For deficient gastric secretion producing fermentation and eructation, one tablet every two hours.

NUX & PEPSIN COMP.....	{ Nux Vom. Po....1-100 gr. }	1 75
	{ Pepsin Pure.....1-4 gr. }	
	{ Bismuth Subnit....1 gr. }	
	{ Calomel.....1-4 gr. }	

One or two after meals, for chronic gastritis, nausea and pain after eating.

NUX AND PHOSPHORUS.....	{ Ext. Nux Vom....1-4 gr. }	1 00
	{ Phosphorus.....1-50 gr. }	

One, three or four times a day, for headache and neuralgia depending upon intestinal dyspepsia.

NUX & SULPHUR.....	{ Nux Vomica Po.1-100 gr. }	75
	{ Sulphur.....1-100 gr. }	

Two or three after a meal, as an aid to digestion.

NUX & SULPHUR COMP.....	{ Nux Vomica Po.1-20 gr. }	75
	{ Sulphur.....1-10 gr. }	
	{ Acid Arsenous...1-100 gr. }	

One after meals, for chronic skin diseases, anæmia, and indigestion.

PEPPERMINT OIL	1-20 gr.....	75
One to two, dissolved in hot water on the end of a spoon, for a carminative in infantile colic.		
PEPSIN PURE	1-20 gr.....	75
Give one after each nursing to an infant with regurgitation and colic.		
PEPSIN PURE	1-10 gr.....	75
One after eating for dyspepsia of children; take whole or crushed in a little water.		
PEPSIN PURE	1-4 gr.....	1 00
One to four tablets after meals, for tardy and incomplete stomach digestion.		
PEPSIN PURE	1 gr.....	2 25
One, three times a day, after eating, for enfeebled digestion.		
PEPSIN PURE	3 gr.....	5 00
A tablet after eating, for dyspepsia.		
PEPSIN COMP	{ Pepsin Pure.....1-4 gr. }	1 25
(Dr. Still.)	{ Ext. Nux Vom.....1-32 gr. }	
	{ Ipecac Powd.....1-16 gr. }	
One to two tablets directly after meals, repeated every half hour, if necessary, for three doses; as a tonic and aid to digestion.		
PEPSIN & BISMUTH	{ Pepsin Pure.....1-4 gr. }	1 75
Antidyspeptic.	{ Bismuth Subnit.....1 gr. }	
One or two at a time, for functional disorders of the stomach.		
PEPSIN, BISMUTH & CHARCOAL	{ Pepsin Conc.....2 gr. }	4 50
	{ Bismuth Subnit.....2 gr. }	
Absorbent stomachic and digestive.	{ Charcoal.....2 gr. }	
A tablet twenty to thirty minutes after meals, for delayed and imperfect digestion with pyrosis and eructation.		
PEPSIN, BISMUTH AND GINGER	{ Pepsin Sacch.....2 gr. }	2 85
	{ Bismuth Subnit.....3 gr. }	
	{ Ginger.....1 gr. }	
One, three or four times a day, for dyspepsia and flatulence of sedentary life.		
PEPSIN BISMUTH & STRYCHNINE	{ Pepsin Pure.....1 gr. }	3 25
	{ Bismuth Subnit.....2 gr. }	
	{ Strychnia Sulph.....1-60 gr. }	
Give one after meals and one at night, decreasing to two a day as symptoms improve; for atonic dyspepsia.		
PEPSIN & CALOMEL	{ Pepsin Pure.....1-4 gr. }	1 85
	{ Bismuth Subnit.....1 gr. }	
	{ Nux Vomica.....1-100 gr. }	
	{ Calomel.....1-40 gr. }	
Three to six a day, as a stomachic tonic and intestinal and hepatic stimulant, for chronic dyspepsia with constipation.		
PEPSIN & CHARCOAL	{ Pepsin Pure.....1-4 gr. }	1 50
	{ Carbo Veg.....1-4 gr. }	
One every hour or two, for flatulent dyspepsia		
PEPSIN & PANCREATIN	{ Pepsin Pure.....1-4 gr. }	2 00
Digestive.	{ Pancreatin.....1-4 gr. }	
One to three after meals, for incomplete and tardy digestion, inability to digest oils and fats, and in the treatment of anæmia and emaciation.		
PEPSIN SACCHARATED	1 gr.....	1 10
One every two hours with some nutriment, in the forced nutrition of consumption and during convalescence.		
PEPSIN SACCHARATED	5 grs.....	1 75
A tablet after meals, as an aid to stomach digestion.		

PHENACETINE	1 gr.....	5 00
One every two hours for a child ; for fever with pain and restlessness. Five at a dose, repeated as necessary, for neuralgia, rheumatism and headache.		
PHENACETINE	2 grs.....	9 00
Analgesic and antipyretic.	Per 100.....	90
One every two hours for migraine ; to quiet restlessness and to reduce fever		
PHENACETINE	3 grs.....	12 50
	Per 100.....	1 25
One every hour, for the fever and pain of acute rheumatism.		
PHENACETINE	4 grs.....	15 50
	Per 100.....	1 00
One or two at a dose, repeated as necessary, to depress temperature and quiet pain and restlessness, in fever and rheumatism.		
PHENACETINE	5 grs.....	18 50
	Per 100.....	1 85
One or two at a dose, repeated as necessary, to depress temperature and quiet pain and restlessness, in fever and rheumatism.		
PHENACETINE COMP.	{ Antipyrine.....2 gr. Phenacetine.....2 gr. Caffeine Cit.....1 gr. Sugar of Milk.....1 gr. Per 100.....	2 50
One to four, according to the height of the temperature, as an antipyretic in all fevers ; with analgesic and diuretic secondary effects.		
PHOSPHATES	{ Iron Phosphate.....1 gr. Quinine Phosphate.....1 gr. Strych. Phos.....1-60 gr. Tonic.	2 75
One three or four times a day as a general tonic and antimalarial ; decrease the dose if ringing in the ears and headache occurs.		
PICROTOXIN	1-80 gr.....	1 10
One every four to six hours, for spinal irritation, petit mal and hyperidrosis.		
PICROTOXIN	1-30 gr.....	1 50
One at night, to control the sweating in phthisis and rheumatism.		
PILOCARPINE MUR	1-100 gr.....	2 25
One every hour, for pleurisy and other serous inflammations.		
PILOCARPINE MUR	1-50 gr.....	4 50
Give a tablet three to six times a day, to increase the flow of saliva and urine.		
PILOCARPINE MUR	1-20 gr.....	10 00
One to three a day, as a sialogogue, diuretic and diaphoretic.		
PILOCARPINE MUR	1-10 gr.....	20 00
One tablet, for a diaphoretic and diuretic action, repeated with care after three or four hours.		
PINK TABLETS (Sugar of Milk)		50
PLUMMER	1-1000 gr.....	75
One to three, morning and night, as an alterative, for a child.		
PLUMMER	1-100 gr.....	75
One or two every two hours, as an hepatic stimulant to the liver, and to rouse glandular and lymphatic action.		
PLUMMER	1-10 gr.....	75
One every two hours.		

When ordering or prescribing

PODOPHYLLIN	1-100 gr.....	75
One every hour for children, until effect is produced, for constipation with clay colored fæces.		
PODOPHYLLIN	1-40 gr.....	75
One every hour till the bowels move freely, then one after meals; for chronic hepatic atony.		
PODOPHYLLIN	1-30 gr.....	75
Two or three at night and three during the next day, to stimulate the excretion of bile in chronic constipation.		
PODOPHYLLIN	1-10 gr.....	75
One every hour or two, for bilious headache; then two at night for several days.		
PODOPHYLLIN	1-8 gr.....	75
One after meals for biliousness and beginning jaundice.		
PODOPHYLLIN	1-4 gr.....	75
One to two at night, and one every three hours next day until a cathartic effect is produced, for biliousness and constipation.		
PODOPHYLLIN	1-2 gr.....	1 10
A tablet at night, for biliousness and habitual headache.		
PODOPHYLLIN COMP.	{ Podophyllin.....1-2 gr. Ext. Nux Vom.....1-10 gr. Ext. Hyoscyam.....1-8 gr. }	1 50
Tonic laxative.		
One or two at bedtime, for atonic conditions of the intestine, and as a stimulant to the liver.		
PODOPHYLLIN AND COLO- CYNTH COMP.	{ Podophyllin.....1-5 gr. Ext. Colocyn.Co. 1 1-2 gr. Ext. Jalap.....1 gr. Ext. Hyoscyamus. 1-5 gr. }	3 25
Cathartic.		
One every hour until effect is produced, as a hydragogue cathartic; one at night in habitual constipation, especially of rheumatic subjects.		
PODOPHYLLIN & LEPTAN- DRIN	{ Podophyllin.....1-4 gr. Leptandrin.....1-2 gr. }	1 50
Hepatic.		
From one to four at night, or divided through the day, for strong hepatic stimulation in functional disorders of the liver.		
POTASS. ARSENITIS	1-5000 gr.....	75
Alterative.		
One tablet to a child, repeated every hour, as a tonic and for skin diseases, also for some kinds of diarrhœa. One tablet equals one-fiftieth minim of Fowler's solution.		
POTASS. ARSENITIS	1-1000 gr.....	75
One tablet every two hours for diseases of the skin, chronic diarrhœa, neuralgia, etc. One tablet equals one-tenth minim of Fowler's solution.		
POTASS. ARSENITIS	1-500 gr.....	75
One every three hours for tonic, also for skin disease; dose may be slowly increased. One tablet equals one-fifth minim of Fowler's solution.		
POTASS. ARSENITIS	1-100 gr.....	75
One tablet equals one minim of Fowler's solution; used in skin diseases, useful also in coryza and some forms of diarrhœa.		
POTASS. ARSENITIS	1-50 gr.....	75
One after meals, increased slowly, for anæmia, chorea and chronic skin diseases. One tablet equals two minims of Fowler's solution.		
POTASS. ARSENITIS	1-40 gr.....	75
One three times a day for chronic rheumatism, malarial poisoning and as an alterative. One tablet equals two and a half minims of Fowler's solution.		

POTASS. ARSENITIS	1-30 gr.....	75
A tablet, three or four times a day for chronic eczema, bronchitis, rheumatism, malarial headache and anæmia. One tablet equals three minims of Fowler's solution.		
POTASS. ARSENITIS	1-25 gr.....	75
One after meals, for pallor and emaciation, asthma and chronic bronchitis; increase slowly as tolerance is established. One tablet equals four minims of Fowler's solution.		
POTASS. ARSENITIS	1-20 gr.....	75
One tablet after meals as a general tonic, in obscure malarial disorders and for acne and eczema. One tablet equals five minims of Fowler's solution.		
POTASS. BICARBONATE	5 grs.....	1 15
Three or four daily, for the uric acid diathesis, to hasten the elimination of urates.		
POTASS. BICARBONATE	10 grs.....	1 50
A tablet three times a day, for sub-acute and chronic rheumatism, gout, and uric acid excess.		
POTASS. BICHROMATE	1-100 gr.....	75
One on the tongue every two hours, for sore throat and catarrhal affections of the lower air passages.		
POTASS. BICHROMATE	1-50 gr.....	75
One on the tongue every two hours, for sore throat and catarrhal affections of the lower air passages.		
POTASS. BICHROMATE	1-10 gr.....	75
One every three hours for catarrhal affections of the lower air passages and as a remedy in catarrhal inflammation of the bowels.		
POTASS. BROMIDE	5 grs.....	1 15
Sedative.		
A tablet every three hours, to induce sleep; one after meals for hypochondria, and chorea; in gradually increasing doses for hysteria, convulsive epilepsy, and threatened abortion.		
POTASS. BROMIDE	10 grs.....	1 00
One to three dissolved in water, repeated if necessary in one hour, or one, three times a day, for nervousness and sleeplessness.		
POTASS. CHLORATE	1-2 gr.....	75
One every half hour for irritation of the throat.		
POTASS. CHLORATE	1 gr.....	75
One every half hour for irritation of the throat.		
POTASS. CHLORATE	3 grs.....	75
One every hour for six doses, allowed to dissolve slowly in the mouth, for hoarseness, tickling and irritation in the throat.		
POTASS. CHLORATE	5 grs.....	75
Throat.		
Dissolve slowly in the mouth every two or three hours, for sore throat, canker and bad breath.		
POTASS. CHLORATE	5 grs.....	75
(With Oil Wintergreen.)		
Sore throat.		
A tablet dissolved on the tongue every two hours, for sore throat and slight colds, and vocal fatigue of speakers and singers.		
POTASS. CHLORATE COMP.	{ Potass. Chlorate.... 1 gr. / (Dr. McEwen.) { Ipecac Powd..... 1-20 gr. \	75
One dissolved slowly on the tongue every hour; for cold with sore throat and hoarseness.		

When ordering or prescribing

POTASS. IODIDE 1 gr. 1 50
 Alterative.

One every hour, for syphilitic and strumous diseases of children; may be combined with chalybeates.

POTASS. IODIDE 2 grs. 2 50

One every hour or two, for chronic rheumatism and diseases of the skin; with free use of drinking water.

POTASS. IODIDE 5 grs. 5 00

One, three times a day, increase gradually to four at a dose, for secondary syphilitic diseases; one every three hours with a swallow of water, for asthma.

POTASS. NITRATE 5 grs. 75

One every three hours, as a diuretic and laxative; lessens fever and pulse-rate, and relieves testicular neuralgia.

POTASS. PERMANGANATE 1 gr. 1 00

One every three hours, for acute rheumatism, diphtheria and zymotic diseases. Also one, three times daily as an emmenagogue; also useful in amenorrhœa.

POTASS. PERMANGANATE 2 grs. 1 15

One, three times daily as an emmenagogue. Also one to four to a pint of hot water, as an injection in vaginitis.

PROSTATIC COMP. } Ext. Saw Palmetto. . . 2 gr. {
 } Ext. Nux Vom. . . . 1-8 gr. { 2 75

One night and morning, for irritability and chronic enlargement of the prostate, and loss of power in the bladder

QUININE ARSENATE 1-64 gr. 90

One every three hours, with gradual increase of the dose, for malarial anæmia and cachexia.

QUININE ARSENATE 1-50 gr. 90

One every three hours, increasing by one a week until five are taken, for dumb ague and chronic malaria.

QUININE, ARSENIC AND } Quinine Sulph. . . . 1 gr. {
STRYCHNINE } Acid Arsenous. . . 1-50 gr. { 2 25
 } Strych. Sulph. . . 1-60 gr. {

One after meals, for a general tonic effect in anæmia and malarial infection.

QUININE & ATROPINE CO. } Quinine Bisulph. . . 1 gr. {
 Tonic. } Potass. Arsen. . . 1-200 gr. { 1 50
 } Atropine Sulph. 1-600 gr. {

One every hour as a tonic sedative, to relieve restlessness and irritability; also in the fevers of children as an antipyretic.

QUININE BISULPHATE 1-10 gr. 1 00

One, three times a day, as an antiperiodic and tonic, for a child.

QUININE BISULPHATE 1-4 gr. 1 10

One, half an hour before eating, as a stomachic tonic.

QUININE BISULPHATE 1-2 gr. 1 25

A tablet three or more times a day, often combined with iron to advantage, as a general tonic.

QUININE BISULPHATE 1 gr. 1 50

One after meals as a tonic and antiperiodic; one every hour or two as an antipyretic in children's fevers.

QUININE BISULPHATE 2 grs. 2 25

One after meals to strengthen the pulse, increase muscular force, and improve the tone of the nervous system.

QUININE BISULPHATE 3 grs. 3 00

One every two hours, with free use of drinking water, as an antipyretic for coughs and fevers in childhood.

QUININE BISULPHATE4 grs.....	4 00
Per 100.....	50

One every hour to four doses, unless relieved sooner, for migraine and malarial hemicrania; one every three hours as a malarial tonic.

QUININE BISULPHATE5 grs.....	5 00
Per 100.....	00

A tablet every two or three hours, as an antipyretic in fever; for chills, and fever, a tablet, at six, four, and two hours before the chill.

QUININE & CHOCOLATE	{ Quinine Tannate. .1 gr. / Tonic for children. { Chocolate..... .50 gr. }	3 75
-------------------------------------	--	------

One to three at a time, as an agreeable tonic for children.

QUININE COMP.	{ Quinine Sulph.....1 gr. / Acid Arsenous.....1-40 gr. / Iron Powd..... .1 gr. / Strych. Sulph.....1-40 gr. / Oil Bl'k Pepper...1-3 gr. }	2 25
General tonic.		

One or two after meals, as a general tonic in enfeebled and anæmic conditions, with or without a malarial element.

QUININE, DIGITALIS COMP. ..	{ Quinine Sulph....1-2 gr. / Strych. Sulph....1-100 gr. / Digitalis Powd...1-2 gr. / Iron Powd.....1-2 gr. }	1 50
(Dr. Fowler.)		
Heart and general tonic.		

A tablet three or four times a day, for anæmia, feeble or irregular heart action, and nervous exhaustion.

QUININE & HYOSCYAMUS CO	{ Morph. Sulph....1-40 gr. / Ext. Hyoscyam...1-10 gr. / Quinine Sulph.....1 gr. }	2 50
Sedative, tonic.		

From one to four a day, for the irritability of exhaustion, as a general tonic and sedative.

QUININE, IODINE COMP.	{ Quinine Sulph.....2 gr. / Acid Arsen.....1-40 gr. / Iodine (Free)1-12 gr. }	3 25
(Dr. Duncan.)		
Alterative.		

QUININE, IPECAC COMP.	{ Quinine Sulph.....1 gr. / Ipecac Powd.....1-8 gr. / Iron Powd.....1-1-2 gr. / Strych. Sulph.....1-40 gr. / Acid Arsenous...1-40 gr. }	1 85
Corrective tonic.		

A tablet three times a day, for convalescence from wasting disease, anæmia and chlorosis, and chronic dyspepsia.

QUININE, IRON & ARSENIC ..	{ Quinine Sulph...1-2 gr. / Iron Powder.....1-2 gr. / Acid Arsenous...1-120 gr. }	1 50
Haematic.		

One every three or four hours, increasing gradually, for simple anæmia and all enfeebled conditions with pallor.

QUININE & LICORICE	{ Quinine Sulph....1-10 gr. / Ext. Licorice.....1-10 gr. }	1 10
-------------------------------------	--	------

One after meals, for children as a tonic and regulator.

QUININE SULPHATE	1-10 gr.....	95
-------------------------------	--------------	----

A tablet three or four times a day, to strengthen the pulse and invigorate the nervous system.

QUININE SULPHATE	1-4 gr.....	1 10
-------------------------------	-------------	------

Three or four a day, for anæmia and during convalescence; best combined with iron.

QUININE SULPHATE	1-2 gr.....	1 25
-------------------------------	-------------	------

One every hour, as a febrifuge and antiperiodic for children, stop if headache and tinnitus aurium occur.

When ordering or prescribing

QUININE SULPHATE1 gr.....	1 50
One every hour, in acute malarial fevers; when symptoms begin to improve, decrease to every three hours, and then to three times a day; one after meals, as a general tonic.	
QUININE SULPHATE2 grs.....	2 25
One, three or four times a day after meals, as a tonic.	
QUININE SULPHATE3 grs.....	3 00
One for tonic three times a day—two to three for breaking up a cold, generally given with five to ten grains of Dover's powder.	
QUININE SULPHATE4 grs.....	4 00
A tablet three or four times a day as a tonic; one every two hours for neuralgia.	
QUININE SULPHATE5 grs.....	5 00
A tablet at six, four, and two hours before the chill in malarial fever; one or two at night, followed by another next morning, generally combined with Dover's Powder, for recent colds.	
QUININE TANNATE COMP. { Quinine Tannate....1 gr. {	3 00
{ Bismuth Subnit....1 gr. {	
{ Opium Powd.....1-4 gr. }	
One every half hour after meals, for three doses, in malarial dyspepsia and chronic catarrhal gastritis.	
REALGAR , (see Arsen. Red Sul.)	
RHEUMATIC { Acid Salicylic.....3 gr. }	2 50
(Dr. J. F. Reilley.) { Wine Colchicum...1 min. }	
{ Guaiac Res.....3 gr. }	
One every two hours until pain is relieved.	
RHEUMATIC No. 2 { Sodium Salicy....2 1-2 gr. }	2 35
{ Fl.Ex.Case.Sag.2 1-2 min. }	
Two every three or four hours, for rheumatism.	
RHEUMATISM { Sodium Salicylate...5 gr. }	5 00
(Dr. Liggett.) { Colchicin.....1-120 gr. }	
{ Codeine Sulph....1-16 gr. }	
One tablet every three hours, until rheumatic pain ceases and a laxative action is obtained.	
RHINITIS , (1/2 strength.)... { Camphor.....1-4 gr. }	1 50
(Dr. Lincoln.) { Fl. Ext. Bellad...1-8 min. }	
{ Quinine Sulph....1-4 gr. }	
One every half hour for four doses, then every hour until throat feels dry; afterwards every three hours until a day after the acute coryza disappears.	
RHINITIS , Full Strength..... { Camphor.....1-2 gr. }	2 00
{ Fl. Ext. Bellad...1-4 min. }	
{ Quinine Sulph....1-2 gr. }	
One every hour for four doses, then every hour and a half until throat feels dry; afterwards every three hours until a day after acute coryza disappears.	
RHUBARB1-2 gr.....	75
One every two hours, for a tonic to the intestinal canal, and mild astringent laxative.	
RHUBARB1 gr.....	75
One every hour until the bowels move, for faecal accumulation; one after meals, repeated in half an hour, for gastric and duodenal catarrh.	
RHUBARB & BISMUTH CO. { Rhubarb Powd....2 gr. }	2 25
Laxative and stomachic. { Ginger Powd....1-2 gr. }	
{ Sodium Bicarb....1 gr. }	
{ Bismuth Subnit....2 gr. }	
One or two at night, as a laxative in cases of dyspepsia and sluggish bowels.	

RHUBARB COMP. U.S.P......1-2 gr..... 75

One after each meal or two at night, as a laxative; increase for purgative effect.

RHUBARB & GINGER CO. ...	{ Rhubarb Powd. .1 1-4 gr. }	
Laxative and stomachic.	{ Sodium Bicarb. .1 1-4 gr. }	
	{ Ginger Powd. . .2 1-2 gr. }	1 60
	{ Cardamom Powd. .5-8 gr. }	
	{ Oil Peppermint. .1-20 gr. }	

A tablet after meals, for dyspepsia, flatulence and constipation.

RHUBARB & HYOSCYAMUS CO.	{ Rhubarb Powd.3 gr. }	
Alterative and laxative.	{ Calomel.1 gr. }	2 00
	{ Ext. Hyoscyamus. .1 gr. }	

A tablet in the morning and one at night, for constipation with collection of mucus in the lower bowel, and dry stools, with flatulence and headache.

RHUBARB & IPECAC	{ Rhubarb Powd. .1-4 gr. }	
(Dr. Hitchcock.)	{ Ipecac Powd.1-10 gr. }	75
Children's laxative.		

One a dose; four to eight daily, as needed in habitual duodenal torpidity, with lack of bile in the passages.

RHUBARB & IPECAC CO.	{ Rhubarb Powd.1 gr. }	
(Roosevelt Hospital)	{ Sodium Bicarb.5 gr. }	1 10
Alkaline laxative & stomachic	{ Ipecac Powd.1-8 gr. }	
	{ Oil Peppermint. 1-20 min. }	

One after each meal, for indigestion with constipation.

RHUBARB & IPECAC COMP.	{ Rhubarb Powd.2 gr. }	
No. 2.....	{ Sodium Bicarb.5 gr. }	1 50
(Roosevelt Hospital.)	{ Ipecac Powd.1-8 gr. }	
	{ Oil Peppermint. 1-20 min. }	

One after each meal, for indigestion with constipation.

RHUBARB & IPECAC COMP.	{ Rhubarb Powd.2 gr. }	
No. 3.....	{ Ipecac Powd.1-4 gr. }	1 60
(Roosevelt Hospital.)	{ Sodium Bicarb.5 gr. }	
	{ Tr. Nux Vomica. .5 min. }	
	{ Oil Peppermint. .1-20 gr. }	

One after each meal, for indigestion with constipation.

RHUBARB & MAGNESIA	{ Rhubarb Powd.2 gr. }	1 25
	{ Magnesia.2 gr. }	

One every three hours, for skin diseases dependent upon disordered digestion, and as an alkaline corrective for summer dyspepsia.

RHUBARB & SODA	{ Rhubarb Powd. .1 1-2 gr. }	
Alkaline laxative.	{ Sodium Bicarb. .1 1-2 gr. }	1 10
	{ Oil Peppermint. .1-7 min. }	

One for an infant, two for a child, two to five at night for an adult, as a laxative; may be repeated every three hours next day if necessary.

RHUBARB & SODA No. 2	{ Rhubarb.3 gr. }	
	{ Sodium Bicarb.3 gr. }	1 65
	{ Oil Peppermint. .1-4 min. }	

One to two as a laxative, may be repeated every three hours next day if necessary.

RHUBARB AND SODIUM SALICYLATE CO.	{ Rhubarb Powd. .1 1-2 gr. }	
Antiseptic and laxative.	{ Sodium Bicarb. .1 1-2 gr. }	3 25
	{ Ipecac Powd.1-5 gr. }	
	{ Sodium Salicylate. 5 gr. }	
	{ Oil Peppermint. .1-7 min. }	

A tablet three times a day, to prevent septic conditions of the intestine, and as a laxative.

SACCHARATED CALOMEL	{ Calomel.....1 gr. }	1 15
	{ Sodium Bicarb.....2 gr. }	
	{ Sugar.....3 gr. }	

One every three hours until cathartic action is obtained, for a hepatic stimulant and regulator.

SACCHARIN1-2 gr.....	3 25
-----------------------------------	------

Substitute for sugar.

One or two will sweeten a cup of coffee or tea. To be employed instead of sugar in diabetic and dyspeptic conditions.

SACCHARIN1 gr.....	5 00
---------------------------------	------

Substitute for sugar.

One will sweeten a cup of coffee or tea. Use as a substitute for sugar in diabetes, rheumatism and other conditions.

SACCHARIN AND SODIUM BICARBONATE	{ Saccharin.....1-2 gr. }	3 25
	{ Sodium Bicarb.....q. s. }	

One as necessary to take the place of sugar in diabetic diet. One or two will sweeten a cup of coffee or tea.

SALICIN1 gr.....	1 50
-------------------------------	------

Tonic.

A tablet three times a day, as a tonic and sedative; every hour to control fever in children.

SALICIN1-2 gr.....	2 50
---------------------------------	------

One every two or three hours as a febrifuge and antiperiodic; causes no stomach or nervous irritation and may thus be used where quinine disagrees.

SALICIN5 grs.....	3 50
--------------------------------	------

Tonic.

One, three times a day, to cut short severe intermittent fever.

SALICYLATE COMP.	{ Sodium Salicyl.....1-2 gr. }	2 65
Antipyretic.	{ Acetanilid.....5 gr. }	

One or two at the first dose, followed by one every three hours if necessary; for rheumatism with pain and high temperature.

SALINE INFUSION Dr. Geo. Ryerson Fowler,	{ Sodium Chlor.....1-2-3 gr. }	2 00
	{ Sodium Sulphate.....1 gr. }	
	{ Sodium Carb.....2-3 gr. }	
	{ Sodium Phos.....1-3 gr. }	
	{ Magnesia Phos.....1-2-3 gr. }	

A combination of the soluble salts of the blood in the proportions in which they exist in the body. Use one tablet to a pint of water previously sterilized by boiling, to wash away chemical antiseptic agents from wounds after irrigation, or as a saline infusion in cases of excessive hemorrhage, shock, etc.

SALOL1-10 gr.....	75
--------------------------------	----

Intestinal antiseptic.

One every hour, as an intestinal antiseptic for diarrhœa and dysentery of children.

SALOL1-2 gr.....	1 35
-------------------------------	------

A tablet every hour or two, as an intestinal antiseptic for diarrhœa and dysentery of children.

SALOL1 gr.....	1 70
-----------------------------	------

One every hour, for inflammation of the bladder and urethra, combined with Sandalwood (2 drop) Tablet; for catheter fever, fermentation of urine in the bladder, and gonorrhœa.

SALOL2 1-2 grs.....	3 00
----------------------------------	------

One every two hours, as an intestinal antiseptic, and one every hour for acute rheumatism.

SALOL3 grs.....	3 65
------------------------------	------

A tablet three or four times a day, as an intestinal antiseptic.

SALOL	5 grs.....	5 50
One to four tablets every two or three hours, as an antipyretic and intestinal antiseptic.		
SALOL COMP	{ Salol.....3 gr. / Antifebrin.....2 gr. }	4 50
One or two every hour, for high temperature, pain and restlessness.		
SALOL COMP. No. 2	{ Salol.....1 gr. / Bismuth Subnit.....1 gr. / Pulv. Opii.....1-40 gr. / Pulv. Camphor...1-20 gr. }	2 50
Give one every hour for four doses, then every two hours; for fermentative diarrhoea of children, with pain.		
SALOL AND CASCARA SAG- RADA	{ Salol.....5 gr. / Ext. Cascara Sag...1 gr. }	7 00
Antiseptic laxative.		
One to four through the day to control intestinal sepsis and produce natural evacuations.		
SALOL AND PHENACETINE {	Salol.....2 1-2 gr. / Phenacetine.....2 1-2 gr. / Per 100.....	1 50
Analgesic and antiseptic		
One every half hour for three doses, then every two hours, for fever and neuralgia.		
SALOL AND PHENACETINE {	Salol.....1 1-4 gr. / Phenacetine.....1 1-4 gr. / Per 100.....	85
No. 2		
One every hour for four or five doses, then every two hours; for fever, with pain and neuralgia.		
SANGUINARIA & CODEINE {	Tr. Sanguinaria...1 min. / Codeine.....1-25 gr. }	1 30
One every hour, as a sedative and expectorant in laryngitis, bronchitis and whooping cough; stop as symptoms improve.		
SANTONIN	1-10 gr.....	1 00
Vermifuge.		
One to four tablets every two hours, followed by a cathartic; as a vermifuge for children.		
SANTONIN	1-2 gr.....	1 25
One or two, according to age, every two hours or oftener, followed by a cathartic; as a vermifuge.		
SANTONIN	1 gr.....	2 00
One every three hours for not more than six doses, followed by a cathartic; as a vermifuge for children.		
SANTONIN & CALOMEL	{ Santonin.....1-2 gr. / Calomel.....1-2 gr. }	1 65
Vermifuge.		
One to two, according to age, every two hours until cathartic action is produced; as a vermifuge for children.		
SANTONIN AND CALOMEL {	Santonin.....1 gr. / Calomel.....1 gr. }	2 10
No. 2		
One every three hours until cathartic action is produced, or not more than four taken; as a vermifuge for children.		
SANTONIN & PODOPHYLLIN {	Santonin.....1-2 gr. / (Dr. Sidney.) Podophyllin.....1-20 gr. }	1 50
One every hour until the bowels move, for round worms in children; lessen the dose for the very young.		
SCIATICA	{ Tr. Aconite.....3-4 min. / Tr. Belladonna...3-4 min. / Tr. Colchicum....3-4 min. / Tr. Cimicifuga...3-4 min. }	1 10
(Dr. Jno. T. Metcalf.)		
Two tablets, every three hours for rheumatism and sciatica.		

SEDATIVE (Dr. Brown.)	{ Sodium Brom. 2 1-2 gr. }	2 50
	{ Potass. Brom. 2 1-2 gr. }	
	{ Ammon. Brom. 2 1-2 gr. }	
	{ Tr. Hyoscyamus. . . 5 min. }	
	{ Tr. Cannabis Ind. . 5 min. }	

One dissolved in water, repeated every half hour, for three or four doses at night; for wakefulness due to overwork, delirium and nervous neuralgia; also for irritable bladder and emissions.

SEDATIVE No. 2	{ Ext. Opium 1-4 gr. }	1 65
	{ Ext. Can. Ind. 1-8 gr. }	

One, repeated twice at intervals of an hour if necessary, for pain and restlessness.

SENEGA COMP.	{ Tr. Squills. 1 min. }	1 10
	{ Tr. Senega. 1 min. }	
	{ Tr. Ipecac. 1 min. }	

(Dr. Cleveland.)

Expectorant.

Dissolve six tablets in twelve teaspoonfuls of water, and give a child a teaspoonful every hour; two teaspoonfuls every hour for an adult.

SENEGA COMP. No. 2	{ Ammon. Carb. 2 gr. }	3 25
	{ Ex. Senega U.S.P. 70. 1 gr. }	
	{ Ext. Hyoscyamus. 1-8 gr. }	
	{ Ipecac Powd. 1-8 gr. }	

(Dr. English.)

{ Oil Cinnamon. . . 1-2 min. }

One tablet every one to four hours, followed by a swallow of water. A stimulant expectorant and sedative, indicated in all acute catarrhal diseases of the respiratory tract, as soon as the feverish and acute stage is past; also in chronic troubles of same nature, and especially in bronchitis of the aged.

SICK HEADACHE	{ Irisin. 1-10 gr. }	1 50
	{ Podophyllin. 1-20 gr. }	
	{ Sanguinarin. 1-20 gr. }	
	{ Nux Vomica Po. 1-20 gr. }	
	{ Euonymin. 1-4 gr. }	

(Dr. Hale.)

One, repeated every hour until effect is produced; for sick headache. Used also as a hepatic corrective.

SODA MINT	{ Sodium Bicarb. 5 gr. }	95
	{ Oil Peppermint. 1-4 min. }	
	{ Arom. Spts. Amm. . 2 gr. }	

Alkaline stomachic.

One as needed, for indigestion, flatulence, heartburn, sick headache, sour stomach, and nausea.

SODA MINT & PEPSIN	{ Soda Mint. 1-2 gr. }	1 50
	{ Pepsin. 1-2 gr. }	

One to four after eating, repeated at intervals of half an hour, for gastric distress and dyspepsia.

SODIUM ARSENATE	1-100 gr.	75
------------------------------	----------------	----

Alterative.

One every three or four hours, increasing slowly until five are taken; stop when oedema of the face and eyelids and disturbance of digestion occur; for chronic diseases of the skin.

SODIUM ARSENATE	1-60 gr.	75
------------------------------	---------------	----

One three times a day, as a tonic and antimalarial; stop if gastric symptoms occur.

SODIUM ARSENATE	1-50 gr.	75
------------------------------	---------------	----

One after meals, for chronic eczema, psoriasis, and other skin diseases; increase slowly and stop on occurrence of gastric disorder and puffiness of the face.

SODIUM ARSENATE	1-25 gr.	75
-----------------------------	---------------	----

Begin with one and increase as tolerance is established; stop if the stomach should become deranged; for chorea, epilepsy and malarial neuralgia.

SODIUM ARSENATE	1-16 gr.....	75
One after meals, for chronic rheumatism, skin diseases and malarial cachexia; stop the treatment on the appearance of facial œdema and gastric distress.		
SODIUM ARSENATE	1-10 gr.....	75
A tablet two or three times a day, for chorea, epilepsy, and neuralgia, chronic diseases of the joints and skin, and malarial infection; stop the treatment if symptoms of reaction occur.		
SODIUM BICARBONATE	5 grs.....	75
Alkaline.		
One or two after eating to correct acidity of the stomach, and as an aid to digestion in dyspepsia. May be repeated in half an hour if necessary.		
SODIUM BICARBONATE	10 grs.....	75
A tablet every half hour, for two or three doses, to correct acid dyspepsia.		
SODIUM BICARBONATE CO.	{ Sodium Bicarb.....2 gr. { { Saccharated Pepsin 1 gr. { (Dr. C. G. Rancourt.) { Rhubarb Powd ...1-2 gr. }	1 25
Four tablets at a dose after meals; for a child, put one in half a glass of water and give during the day. Useful in gastric fever with a heavily coated tongue.		
SODIUM BROMIDE	5 grs.....	1 30
Sedative.		
One dissolved in water every fifteen minutes for three doses. May be repeated in two hours, for nervousness, restlessness and sleeplessness.		
SODIUM BROMIDE	10 grs.....	1 75
One every hour for three doses to induce sleep; one, three or four times a day, for nervous irritability.		
SODIUM PHOSPHATE	5 gr.....	1 00
Three at night as a mild saline laxative; one after each meal for duodenal catarrh, and the uric acid diathesis.		
SODIUM SALICYLATE	1-2 gr.....	1 10
Anti-rheumatic, analgesic.		
A tablet every hour in acute rheumatism, as an antipyretic and analgesic.		
SODIUM SALICYLATE	1 gr.....	1 50
One every hour, for rheumatism or sour stomach, or five every three hours, for rheumatism.		
SODIUM SALICYLATE	1 gr.....	1 50
(With Oil Wintergreen.)		
A tablet every hour, for the fever and pain of acute rheumatism diminish as symptoms improve.		
SODIUM SALICYLATE	2 grs.....	1 75
Anti-Rheumatic.		
One every hour or two in acute articular rheumatism and serous inflammations of rheumatic origin; decrease the dose if nausea and vomiting occur.		
SODIUM SALICYLATE	3 grs.....	2 25
One every two hours for three doses, for the headache and neuralgia of rheumatic patients.		
SODIUM SALICYLATE	5 grs.....	3 00
One tablet every two to four hours, for gout and rheumatism.		
SODIUM SALICYLATE CO.	{ Sodium Salicylate...2 gr. { { Cerium Oxalate.....1 gr. }	2 00
A tablet every two hours, for pericarditis and acute rheumatism.		
SODIUM SULPHITE	5 grs.....	1 50
Intestinal antiseptic.		
One, three or four times a day, for intestinal stasis and fermentation.		

SODIUM SULPHOCARB.1 gr.....	1 10
One every hour for five or six doses, then every two hours, for an intestinal antiseptic and to reduce temperature.	
SODIUM SULPHOCARB.2 1-2 grs.....	1 35
One or two every hour, to reduce the temperature, correct the bowels and promote intestinal antiseptics in typhoid fever and gastro-enteritis.	
SORE THROAT	1 50
Diphtheritic.	
One every two hours.	
SORE THROAT No. 2	1 50
Tr. Chlor. Iron... 3 min. } Corros. Sublim... 1-100 gr. } Tr. Aconite..... 1-2 min. }	
Tr. Chlor. Iron... 5 min. } Corrosive Sub... 1-64 gr. }	
For tonsillitis, give one every three hours for children of 6 to 12 years. Two tablets every four hours for patients above 12 years old. For diphtheria, crush and give one every two or three hours.	
SPARTEINE SULPHATE1-100 gr.....	75
Heart tonic.	
A tablet every hour or two for irregular and feeble heart action.	
SPARTEINE SULPHATE1-50 gr.....	75
A tablet every two or three hours, to regulate the heart and increase the flow of urine.	
SPARTEINE SULPHATE1-20 gr.....	1 10
One every three hours, as a diuretic in renal congestion and to control irregular cardiac action.	
SPARTEINE SULPHATE1-10 gr.....	1 60
A tablet three to six times a day, as a diuretic in albuminuria and dropsy.	
SPARTEINE SULPHATE1-4 gr.....	2 50
One every three or four hours to favor the elimination of urine and serous exudation, and to correct feeble, irregular heart action.	
SPONGIA1-10 gr.....	75
Throat	
Give one to four every two or three hours, in recent affections of the throat and trachea.	
STIBIUM, OPIUM AND CAM-	75
PHOR.	
(Dr. Fowler.)	
Sedative and expectorant.	
One every half hour or every hour, upon the earliest intimations of a cold.	
STOMACHIC	1 50
(Dr. North.)	
Pepsin Pure..... 1-4 gr. }	
Ext. Nux Vomica... 1-4 gr. }	
Carbo Veg..... 1-4 gr. }	
Capsicum Powd... 1-4 gr. }	
One after meals, repeated in half an hour, for catarrhal inflammation of the stomach, atonic dyspepsia, acid eructation and flatulence.	
STOMACHIC No. 2	1 50
(Dr. North.)	
Pepsin Pure..... 1-4 gr. }	
Ext. Nux Vom... 1-8 gr. }	
Carbo. Veg..... 1-2 gr. }	
Capsicum Powd... 1-8 gr. }	
One or more after meals, for indigestion and gastric atony.	
STRONTIUM COMP.	4 00
(Dr. Winfield.)	
Strontium Lact... 2 1-2 gr. }	
Sparteine Sulph... 1-24 gr. }	
Sulphur Sublim... 1-4 gr. }	
From six to fifteen tablets during the twenty four hours, according to indications, for gout, lithæmia, renal insufficiency and cardiac weakness.	

STROPHANTHUS COMP	Tr. Digitalis 3 min. }	1 50
Heart tonic.	Tr. Strophanthus . . . 2 min. }	

One every two hours, decreasing to two or three a day, for rapid, feeble and irregular action of the heart.

STROPHANTHUS COMP. No.2	Tr. Digitalis 3 min. }	1 00
	Tr. Strophanthus . . . 2 min. }	
	Nitroglycerin . . . 1-100 gr. }	

A tablet three times a day, for rapid and feeble cardiac action, with pallor of the face and cerebral anæmia, and neuralgia.

STROPHANTHUS COMP. No.3	Tr. Digitalis 3 min. }	1 75
(Dr. Pike.)	Tr. Strophanthus . . . 2 min. }	
	Tr. Nux Vomica . . . 2 min. }	

One midway between meals, as a cardiac and stomachic tonic, for cases of irregular and feeble heart action and a tendency to nervous congestion.

STRYCHNINE ARSENATE . . . 1-100 gr.	75
--	----

A tablet after meals, for general venous stasis of the abdominal organs, and as a nerve tonic.

STRYCHNINE ARSENATE . . . 1-50 gr.	75
---	----

One after meals, as a general tonic ; one every three hours, with care, as a diuretic in passive renal congestion.

STRYCHNINE ARSENATE . . . 1-40 gr.	75
---	----

A tablet three times a day, as a tonic and diuretic.

STRYCHNINE MURIATE . . . 1-100 gr.	1 10
---	------

One every three hours, as a nerve and stomachic tonic.

STRYCHNINE MURIATE . . . 1-60 gr.	1 10
--	------

One every three hours, as a nerve and stomachic tonic.

STRYCHNINE MURIATE . . . 1-30 gr.	1 10
--	------

One every three hours, as a nerve and stomachic tonic.

STRYCHNINE MURIATE . . . 1-20 gr.	1 10
--	------

One, three times a day, as a general nerve tonic.

STRYCHNINE MURIATE CO.	Corrosive Sub. . . 1-20 gr. }	1 10
	Ext. Belladonna . 1-10 gr. }	
	Strychnia Mur. . . 1-40 gr. }	

One night and morning for the correction of atonic conditions of the alimentary canal and consequent reflex disorders.

STRYCHNINE NITRATE . . . 1-100 gr.	1 00
---	------

A tablet three times a day, as a general tonic in neurasthenia and spinal anæmia.

STRYCHNINE NITRATE . . . 1-60 gr.	1 00
--	------

One after meals for amenorrhœa, with anæmia and a tendency to melan cholia.

STRYCHNINE NITRATE . . . 1-40 gr.	1 00
--	------

One after eating, for dyspepsia with anorexia, constipation and epigastric distress.

STRYCHNINE NITRATE . . . 1-30 gr.	1 00
--	------

One every four hours, for vesical atony and dyspepsia ; decrease as symptoms improve.

STRYCHNINE NITRATE . . . 1-20 gr.	1 00
--	------

One tablet for the weakness and nervous depression following excessive use of alcohol or tobacco ; may be repeated in two hours.

STRYCHNINE SULPHATE . . . 1-1000 gr.	75
---	----

One every two hours, as a stomachic and general nerve tonic.

STRYCHNINE SULPHATE . . . 1-500 gr.	75
--	----

One every two hours, as a stomachic and general nerve tonic.

When ordering or prescribing

STRYCHNINE SULPHATE . . . 1-200 gr.	75
One every three hours, as a stomachic and general nerve tonic.	
STRYCHNINE SULPHATE . . . 1-100 gr.	75
One, three times a day, as a stomachic and general nerve tonic.	
STRYCHNINE SULPHATE . . . 1-60 gr.	75
One after each meal, for a nerve tonic and aid to digestion.	
STRYCHNINE SULPHATE . . . 1-50 gr.	75
One after each meal, as a nerve tonic and aid to digestion.	
STRYCHNINE SULPHATE . . . 1-40 gr.	75
One after each meal, as a nerve tonic.	
STRYCHNINE SULPHATE . . . 1-30 gr.	75
One after each meal, as a nerve tonic.	
STRYCHNINE SULPHATE . . . 1-20 gr.	75
One after each meal, as a nerve tonic.	
STRYCHNINE SULPHATE . . . 1-16 gr.	75
STRYCHNINE SULPHATE . . . 1-10 gr.	95
STRYCHNINE VALER 1-100 gr.	I 10
One, three times a day, for a general nerve tonic.	
STRYCHNINE VALER 1-60 gr.	I 10
One, three times a day, for a general nerve tonic.	
STRYCHNINE VALER 1-50 gr.	I 10
A tablet night and morning, increasing to three or four a day if well tolerated, for nervous diseases and prostration, and as a tonic aphrodisiac.	
SULPHONAL 2 grs.	11 25
Hypnotic.	
Per 100.	I 15
One every hour for three doses, for restlessness and insomnia.	
SULPHONAL 3 grs.	16 90
Per 100.	I 70
A tablet every hour or two for three or four doses, for insomnia.	
SULPHONAL 5 grs.	26 25
Per 100.	2 65
One every hour or two, as a hypnotic.	
SULPHONAL 8 grs.	40 00
Per 100.	4 00
One or two, one hour before retiring, repeated every hour for two doses if necessary.	
SULPHONAL 15 grs.	70 00
Per 100.	7 00
One an hour before bedtime for insomnia; repeat at bedtime, then another in an hour if necessary.	
SULPHONAL AND HYOSCIN (Sulphon. 5 gr.)	
Sedative and hypnotic. (Hyoscin Hydro. 1-200 gr.)	
Per 100.	3 75
One, three times a day, or every three hours at night, if necessary, as a sedative and hypnotic.	
SULPHUR 1-100 gr.	75
One every two hours, as a laxative for an infant, and in the chronic stage of pertussis.	
SULPHUR 1-50 gr.	75
A tablet three or four times a day, with local applications, for subacute and chronic skin diseases of children.	
SULPHUR 1-10 gr.	75
One every three hours, for persistent bronchitis after whooping cough.	

SULPHUR	1-2 gr.....	75
One every three hours for chronic catarrh, rheumatism and skin diseases.		
SULPHUR	1 gr.....	75
One or two after meals, for acne, pityriasis, and other skin diseases, and as a laxative for children.		
SULPHUR COMP.	{ Sulphur.....5 gr. / Cream Tartar.....1 gr. }	1 00
One or two at night, as a mild stimulating laxative.		
SULPHUR COMP. No. 2	{ Sulphur.....2 1-2 gr. / Cream Tartar...2 1-2 gr. }	1 00
One after meals and two at night, for hæmorrhoids, and as a laxative in pregnancy.		
SULPHUR COMP. No. 3	{ Sulphur.....5 gr. / Cream Tartar.....2 gr. / Ext. Ipecac.....1-100 gr. / Ext. Capsicum...1-500 gr. / Acid Arsen....1-1000 gr. / Calcium Sulph...1-8 gr. }	1 50
Two to four a day, for chronic eczema, pityriasis, acne, and other obstinate cutaneous diseases.		
SULPHUR & IPECAC	{ Sulphur.....1-10 gr. / Ipecac Powd....1-10 gr. }	75
(Dr. Gilbert.)		
One every three hours, in gastric derangements and as an aid to digestion.		
SUN CHOLERA TABLETS ..	{ Tr. Opii.....3 min. / Tr. Rhei.....3 min. / Tr. Capsici.....3 min. / Spts. Camphoræ...3 min. / Spts. Ment. Pip..3 min. }	2 75
Each tablet represents 15 minims of Sun Cholera Mixture.		
Give two tablets at the first dose; then one after each loose movement, for six or eight doses if necessary.		
SUN CHOLERA TABLETS ...	{ Tr. Opii.....2 min. / Tr. Rhei.....2 min. / Tr. Capsici.....2 min. / Spts. Camphoræ...2 min. / Spts. Ment. Pip..2 min. }	2 00
No. 2		
Each tablet represents 10 minims of Sun Cholera Mixture.		
Give two tablets at the first dose; then one after each loose movement, for six or eight doses if necessary.		
SYR. SQUILLS COMP.	5 min.....	1 50
One every three hours, or oftener, for winter cough, asthma, pneumonia and phthisis.		
TARTAR EMETIC	1-100 gr.....	75
One every two hours for a child, in bronchial affections; two for an adult.		
TARTAR EMETIC	1-50 gr.....	75
One every two or three hours for adult, for bronchial affections.		
TARTAR EMETIC	1-20 gr.....	75
A tablet three times a day, for cough, and chronic bronchitis with scanty expectoration.		
TARTAR EMETIC	1-10 gr.....	75
Dissolve one tablet in four tablespoonfuls of water, and of this give a teaspoonful every half hour for an infant with bronchitis.		
TARTAR EMETIC COMP	{ Opium Powd.....1-8 gr. / Tart. Emetic....1-16 gr. / Calomel.....1-2 gr. }	00
Expectorant.		
A tablet three or four times a day, for acute and chronic inflammations of the respiratory tract.		

TARTAR EMETIC AND MORPHIA	{ Tartar Emetic...1-100 gr. / Morph. Sulph...1-100 gr. }	75
One every two hours for cough with fever and restlessness of children.		
TARTAR EMETIC & OPIUM	{ Tartar Emetic...1-100 gr. / Opium Powd....1-50 gr. }	75
Sedative expectorant.		
One every two hours, as a sedative expectorant.		
TERPIN HYDRATE	1 gr.....	1 25
Two tablets, followed by one every two hours, for winter cough.		
TERPIN HYDRATE	2 grs.....	1 75
Winter cough.		
One, repeated every two hours, for winter cough.		
TERPIN HYDRATE	3 grs.....	2 00
One every three hours, for cough.		
TERPIN HYDRATE	5 grs.....	2 35
One every three hours, for persistent cough.		
THIERSCH'S SOLUTION TABLETS	{ Boracic Acid...12 gr. / Salicylic Acid.....2 gr. }	2 50
Dissolve a tablet in four tablespoonfuls of hot water to make Thiersch's solution.		
THROAT	{ Potass. Iodide...1-10 gr. / Salt.....1-10 gr. / Sugar.....1-2 gr. }	1 10
Allow a tablet to dissolve in the throat every half hour, for hoarseness, dryness and tickling.		
THROAT TABLETS	{ Acid Benzoic.....1-5 gr. / Tr. Belladonna...1 min. / Paregoric.....10 min. / Ext. Licorice.....q. s. }	1 50
(Dr. F. J. Quinlan.)		
Dissolve one slowly on the tongue, not oftener than every four hours, for dryness of the throat attended with pain. Used also for the relief of acute and subacute inflammation of the fauces.		
TINCT CINCHONA COMP	10 minim.....	1 75
One tablet for a child, three times a day, as a tonic and febrifuge.		
TINCT CINCHONA COMP	1 teaspoonful.....	5 00
One tablet three or four times a day, for tonic.		
TINCT. THUJA	5 minims.....	1 50
One tablet every three hours, for cystitis.		
TONIC	{ Iron Pyrophos...1-2 gr. / Quinine Sulph....1-2 gr. / Strych. Sulph...1-120 gr. }	1 50
(Dr. Hammond.)		
One to two tablets just after each meal, as a nerve or general tonic in anæmia.		
TONIC No. 3	{ Ferri Pyrophos.....2 gr. / Quinine Sulph....1-4 gr. / Strych. Sulph....1-60 gr. }	1 75
(Dr. Hoopes.)		
A tablet three times a day after wasting disease, and as a hæmætic in anæmia and malarial poisoning.		
TONIC ALTERATIVE	{ Corrosive Sub...1-50 gr. / Strych. Sulph...1-100 gr. / Ipecac Powd....1-20 gr. }	1 00
One or two after meals, as a tonic and alterative.		
TONIC ALTERATIVE No. 2	{ Corrosive Sub...1-30 gr. / Strych. Sulph....1-60 gr. / Iron Powder...1 gr. }	1 10
Dr. E. Jay Fisk.		
One to two after meals for anæmic and cachectic conditions, chronic bronchitis and malaria.		

TONIC CHALYBEATE	Plain or Sugar Coated....	2 00
(Dr. Flint.)	1-2 size.....	1 50

From an analysis of the blood, containing an excess of iron and chloride of sodium.

Two tablets, three times a day after eating. A tonic made from an analysis of the blood; useful in anæmia, Bright's disease and nervous exhaustion.

TONIC CHLORIDES	Iron Chloride.....1 gr. }	1 25
	Merc. Bichlor.....1-4 gr. }	
	Arsenic Chlor.....1-64 gr. }	

One after meals for anæmia, and as a general tonic in convalescence.

TONIC GOUT	Quinine Sulphate...2 gr. }	4 50
(Dr. Robbins.)	Ext. Digitalis.....1-4 gr. }	
	Ext. Colchic. Seed...1 gr. }	

A tablet three times a day, until brisk cathartic action is produced; for gout, rheumatism, lumbago and rheumatic neuralgia,

TONIC LAXATIVE	Ext. Colocy. Co...1-2 gr. }	2 80
(Dr. Miller.)	Ext. Belladonna...1-10 gr. }	
	Quinine Sulph.....1 gr. }	

One to three at night, for chronic atony of the bowels and constipation of malarial and anæmic subjects.

TONIC SYPHILITIC	Mercury Protiod...1-5 gr. }	1 50
(Dr. Feltwell.)	Morph. Sulph....1-24 gr. }	
	Tartar Emetic....1-24 gr. }	

TONIC TREATMENT	Capsicum.....1-60 gr. }	1 25
	Opium Powd....1-15 gr. }	
	Merc. Protiodide..1-4 gr. }	

One or two at a dose, two to three times a day.

TONSILLITIS	Tr. Aconite.....1-5 min. }	1 00
	Tr. Bryonia....1-10 min. }	
	Tr. Belladonna...1-10 min. }	
	Merc. Iod. Red...1-100 gr. }	

Dissolve one on the tongue every twenty minutes for first hour, then every thirty minutes for the second hour, after which, one every hour, avoiding all acids or fruit; for sore throat from cold, also in mild cases of diphtheria, and scarlatinal sore throat.

TONSILLITIS No. 3	Corros. Sublim....1-96 gr. }	1 50
(Dr. Bierhoff.)	Tr. Chloride Iron...3 min. }	
	Oil Cinnamon.....q. s. }	

One dissolved in water every hour, for diphtheria or tonsillitis.

TRIONAL	5 grs.....	30 00
----------------------	------------	-------

One every hour, for three doses in insomnia and delirium, and to replace morphine where it disagrees, or in the cure of the habit.

TRIPLE BROMIDES No. 1 ...	Sodium Brom...2 1-2 gr. }	1 35
Sedative.	Potass. Brom...2 1-2 gr. }	
	Ammon. Brom...2 1-2 gr. }	

One to four tablets dissolved in water, every three hours until the patient is quiet.

TRIPLE BROMIDES No. 2 ..	Sodium Bromide...5 gr. }	2 25
	Potass. Bromide...5 gr. }	
	Ammon. Bromide...5 gr. }	

One or two dissolved in water, three times a day, as a nerve sedative.

TRIPLEX	Aloes.....2 gr. }	1 75
	Pil. Hydrarg.....1 gr. }	
	Podophyllin.....1-4 gr. }	

One for a laxative—two for a cathartic—three for a purge; best taken just before retiring, for biliousness and constipation.

TULLY'S COMP.	{ Tully's Tablets...2 1-2 gr. }	1 70
Sedative and diaphoretic.	{ Calomel.... 1-4 gr. }	
One every two or three hours, as a sedative and diaphoretic, in bronchitis and painful inflammations of the thorax.		
TULLY'S COMP. No. 2.	{ Tully's Powder...2 1-2 gr. }	1 70
	{ Podophyllin.... 1-10 gr. }	
A tablet every two or three hours, for fever, pain and cough, and to induce sleep.		
TULLY'S POWDER	1-10 gr.	75
One every two hours to quiet pain and induce sleep; for an infant.		
TULLY'S POWDER	1-5 gr.	75
A tablet every two hours for painful diarrhoea of children.		
TULLY'S POWDER	1-2 gr.	75
One, repeated in an hour, for painful cough of children.		
TULLY'S POWDER	1 gr.	90
One every hour for three or four doses, as a diaphoretic.		
TULLY'S POWDER	5 grs.	2 25
One tablet repeated in an hour and a half, as a sedative and diaphoretic.		
TULLY'S POWDER	10 grs.	3 25
Anodyne.		
One at night, for insomnia due to pain, and for delirium and coma vigil.		
TULLY'S TABLETS	2 1-2 gr.	1 50
A tablet as often as every two hours, as a sedative and anodyne.		
TURPETH MINERAL	1 gr.	75
Emetic.		
One every half hour for three doses, to produce vomiting; aided by warm water.		
TURPETH MINERAL	2 grs.	1 15
A tablet, repeated in half an hour, to empty an overloaded stomach and for croup.		
UTERINE SEDATIVE	{ Helonin.....1-8 gr. }	2 50
(Dr. R. G. H. Hayes.)	{ Viburnin.....1-8 gr. }	
	{ Caulophyllin....1-4 gr. }	
	{ Mitchellia Rep...1 1-2 gr. }	
One to three tablets every three hours, for membranous or neuralgic dysmenorrhœa, and as a uterine tonic or antispasmodic.		
URETHRITIS No. 1	{ Zinc Acetate.....2 gr. }	1 50
For 2 oz. ($\frac{1}{8}$ pint) bottles.	{ Corrosive Sub....1-5 gr. }	
	Per doz. bottles net.....	2 25
Dissolve a tablet in four tablespoonfuls of hot water and use as an injection as hot as can be borne, after each urination.		
URETHRITIS No. 2	{ Zinc Acetate.....4 gr. }	1 75
For 4 oz. ($\frac{1}{4}$ pint) bottle.	{ Corros. Sub.....2-5 gr. }	
Dissolve a tablet in eight tablespoonfuls of hot water and use as an injection as hot as can be borne, after each urination.		
VAGINAL ASTRINGENT	{ Alum.....3 gr. }	2 00
	{ Zinc Sulph.....2 gr. }	
	{ Acid Tannic.....2 gr. }	
	{ Acid Boracic.....6 gr. }	
	{ Ext. Hydrastis.....1 gr. }	
	(Colorless)	
Dissolve two tablets in a quart of water and use as an injection.		
VERATRIA SULPHATE	1-20 gr.	1 30
A tablet every four hours, for pertussis, and neuralgia.		

VERATRIA SULPHATE.1-10 gr. 2 25

One every three or four hours, for gout, rheumatism, neuralgia and paralysis agitans.

VIBURNUM SEDATIVE. 2 75

(Ext. Viburnum Prun.1 gr.	}	
Ext. Viburnum Op..1 gr.		
Ext. Aletris Far...1-2 gr.		
Ext. Helon. Dioica.1-2 gr.		
Ext. Mitchel. Rep.1-2 gr.		
Caulophyllin.....1-4 gr.		

One to three tablets every three hours, or one tablet every hour, with a swallow of hot water or tea, as a uterine tonic and antispasmodic.

VIBURNUM SEDATIVE No. 2. 1 00

(Fl. Ex. Viburn. Op..1 min.	}	
Camphor.....1-2 gr.		
Opium Powd.1-4 gr.		
Oil Anise.....1-8 min.		

Begin one day before the period, with a tablet night and morning; the next day take three and the next four; for scanty and painful menstruation and ovarian colic.

VIBURNUM SEDATIVE No. 3. 2 00

(Dr. Fisher.)	}	
Black Haw.....3 3-4 gr.		
Golden Seal.....3 3-4 gr.		
Each tablets represents :	}	Jam. Dogwood...1 3-4 gr.)

One after meals for the week before the catamenia, increase if necessary during the flow; for dysmenorrhœa, menorrhagia, and hyperæmia of the pelvic organs, and as an antispasmodic.

WHOOPING COUGH. 4 50

(Antipyrin.....1-2 gr.	}	
Tr. Drosera.....1 min.		
Belladonna.....1-500 gr.		
Ipecac Powd.....1-200 gr.)		

One tablet every hour, for whooping cough.

WHOOPING COUGH No. 2. 1 50

(Acetanilid.....1-2 gr.	}	
Tr. Drosera.....1 min.		
Belladonna.....1-400 gr.		
Ipecac Powd.....1-200 gr.)		

One tablet every hour, for whooping cough.

YELLOW TABLETS (Sugar Milk) 50

ZINC ACETATE.....1 gr. 75

One dissolved in one ounce in water, as an injection.

ZINC COMP. 1 50

(Lead Acetate.....2 gr.	}	
Zinc Acetate.....2 gr.)		

Two dissolved in four ounces of water, for injection.

ZINC PHOSPHIDE.....1-100 gr. 75

Nerve tonic.

One after meals, increasing gradually, for chorea and other spasmodic diseases.

ZINC PHOSPHIDE.....1-50 gr. 75

Three or four a day, increasing slowly, for chorea, epilepsy and neuralgia.

ZINC PHOSPHIDE.....1-25 gr. 75

A tablet every four hours, for hysteria and spasmodic affections of the muscles.

ZINC PHOSPHIDE.....1-10 gr. 75

One three times a day, for chronic nervous disease of an ataxic nature, and for neuralgia.

ZINC PHOSPHIDE.....1-4 gr. 1 10

One after meals, as a tonic to the cerebro-spinal centres, in neurasthenia, hysteria and epilepsy.

CIMICIFUGA.....	1-2 minim, 1 minim....	75
CIMICIFUGA.....	2 minims.....	95
CINCHONA.....	1 minim.....	75
CINCHONA.....	2 minims.....	95
COLCHICUM.....	1 minim.....	75
COLCHICUM.....	2 minims.....	95
CONIUM.....	1 minim.....	75
CONIUM.....	2 minims.....	95
CUBEBS.....	1-2 minim, 1 minim....	75
CUBEBS.....	2 minims.....	95
DIGITALIS.....	1-2 minim, 1 minim....	75
DIGITALIS.....	2 minims.....	95
DIGITALIS.....	5 minims.....	1 00
GELSEMIUM.....	1 minim.....	75
GELSEMIUM.....	2 minims.....	95
GUAIAC.....	1 minim.....	75
GUAIAC.....	2 minims.....	95
HYDRASTIS.....	1 minim.....	75
HYDRASTIS.....	2 minims.....	95
HYOSCYAMUS.....	1 minim.....	75
HYOSCYAMUS.....	2 minims.....	95
HAMAMELIS.....	1 minim.....	75
HAMAMELIS.....	2 minims.....	95
IGNATIA.....	1-2 minim, 1 minim....	75
IGNATIA.....	2 minims.....	95
IPECAC.....	1-2 minim, 1 minim....	75
IPECAC.....	2 minims.....	95
KRAMERIA.....	1 minim.....	75
KRAMERIA.....	2 minims.....	95
NUX VOMICA.....	1-2 minim, 1 minim....	75
NUX VOMICA.....	2 minims.....	95
NUX VOMICA.....	5 minims.....	1 10
OPIUM DEOD.....	1 minim.....	1 35
OPIUM DEOD.....	2 minims.....	1 85
SANGUINARIA.....	1 minim.....	75
SANGUINARIA.....	2 minims.....	95
SQUILLS.....	1 minim.....	75
SQUILLS.....	2 minims.....	95
STRAMONIUM.....	1 minim.....	75
STRAMONIUM.....	2 minims.....	95
STROPHANTHUS.....	1 minim.....	95
STROPHANTHUS.....	2 minims.....	1 25
STROPHANTHUS.....	5 minims.....	1 75
VERATRUM VIRIDE.....	1 minim.....	75
VERATRUM VIRIDE.....	2 minims.....	95
VERATRUM VIR. (Norwood's).....	1 minim.....	1 50
VERATRUM VIR. (Norwood's).....	2 minims.....	3 00

Tablets from Mother Tinctures.

ACONITE.	} 1-4 min., 1-2 min., 3-4 min. }	75
ACONITE.	1 min.	
ACONITE.	2 minims.	1 10
ACTÆA RACEMOSA.	1-2 minim.	75
ACTÆA RACEMOSA.	1 minim.	95
ACTÆA RACEMOSA.	2 minims.	1 10
AILANTHUS.	1 minim.	1 30
ALOE.	1 minim.	75
APIS.	1-2 minim, 1 minim.	75
APOCYNUM.	1 minim.	75
APOCYNUM.	2 minims.	1 10
ARNICA.	1 minim.	95
ASCLEPIAS TUB.	1 minim.	75
ASCLEPIAS TUB.	2 minims.	1 10
BAPTISIA.	1-2 minim, 1 minim.	75
BAPTISIA.	2 minims.	1 10
BELLADONNA.	1-2 minim, 1 minim.	75
BELLADONNA.	2 minims.	1 10
BRYONIA.	1-2 minim, 1 minim.	75
BRYONIA.	2 minims.	1 10
CACTUS.	1 minim.	1 10
CALABAR.	1-2 minim.	75
CALABAR.	1 minim.	95
CALABAR.	2 minims.	1 10
CALENDULA.	1 minim.	75
CANNABIS INDICA.	1 minim.	75
CANNABIS SATIVA.	1 minim.	75
CANTHARIDES.	1-2 minim, 1 minim.	75
CANTHARIDES.	2 minim.	95
CAPSICUM.	1-2 minim, 1 minim.	75
CAPSICUM.	2 minims.	95
CAUSTICUM.	1-2 minim.	75
CAUSTICUM.	1 minim.	95
CAUSTICUM.	2 minims.	1 10
CHAMOMILLA.	1 minim.	75
CHAMOMILLA.	2 minims.	1 10
CHELIDONIUM.	1 minim.	75
CHIMAPHILLA.	1 minim.	95
CHINA.	1 minim.	75
CHINA.	2 minims.	95
CIMICIFUGA.	1-2 minim, 1 minim.	75
CIMICIFUGA.	2 minims.	1 10
CISTUS CAN.	1 minim.	75
CLEMATIS.	1 minim.	75
COCCULUS.	1-2 minim, 1 minim.	75

COCCULUS.....	2 minims.....	1 10
COFFEA.....	1 minim.....	75
COFFEA.....	2 minims.....	1 10
COLCHICUM.....	1-2 minim, 1 minim.....	75
COLCHICUM.....	2 minims.....	1 10
COLLINSONIA.....	1-2 minim, 1 minim.....	75
COLLINSONIA.....	2 minims.....	1 10
COLOCYNTHIS.....	1-2 minim, 1 minim.....	75
COLOCYNTHIS.....	2 minims.....	1 10
CONIUM.....	1 minim.....	75
CONIUM.....	2 minims.....	1 10
CORYDALIS.....	1 minim.....	75
CUBEBS.....	1-2 minim, 1 minim.....	75
CUBEBS.....	2 minims.....	95
DIGITALIS.....	1-2 minim, 1 minim.....	75
DIGITALIS.....	2 minims.....	1 10
DIOSCOREA.....	1-2 minim, 1 minim.....	75
DIOSCOREA.....	2 minims.....	1 10
DROSER.....	1-2 minim, 1 minim.....	75
DROSER.....	2 minims.....	1 10
DULCAMARA.....	1 minim.....	75
EUPHORBIA.....	1 minim.....	75
EUPHRASIA.....	1 minim.....	75
EUONYMUS.....	1 minim.....	75
GELSEMIUM.....	1-2 minim, 1 minim.....	75
GELSEMIUM.....	2 minims.....	1 15
GUAIAC.....	1 minim.....	75
HAMAMELIS.....	1 minim.....	75
HAMAMELIS.....	2 minims.....	95
HYDRASTIS.....	1 minim.....	75
HYDRASTIS.....	2 minims.....	95
HYOSCYAMUS.....	1 minim.....	75
HYOSCYAMUS.....	2 minims.....	1 10
IGNATIA.....	1-2 minim, 1 minim.....	75
IGNATIA.....	2 minims.....	95
IPECAC.....	1-2 minim, 1 minim.....	75
IPECAC.....	2 minims.....	95
IRIS.....	1 minim.....	75
KALMIA.....	1 minim.....	75
LEPTANDRA.....	1 minim.....	75
LOBELIA.....	1 minim.....	75
NUX VOMICA.....	1-2 minim, 1 minim.....	75
NUX VOMICA.....	2 minims.....	95
OPIUM.....	1-2 minim.....	95
OPIUM.....	1 minim.....	1 30
OPIUM.....	2 minims.....	1 85
PHOSPHORUS.....	1 minim.....	75
PHOSPHORUS.....	2 minims.....	95

When ordering or prescribing

PHYTOLACCA.....	1 minim.....	75
PODOPHYLLUM.....	1 minim.....	75
PODOPHYLLUM.....	2 minims.....	1 15
PULSATILLA.....	1-2 minim, 1 minim.....	75
PULSATILLA.....	2 minims.....	1 15
RHUS TOX.....	1 minim.....	75
RHUS TOX.....	2 minims.....	1 15
RUMEX.....	1 minim.....	75
SAMBUCUS.....	1 minim.....	75
SENECIO.....	1 minim.....	75
SPIGELIA.....	1 minim.....	75
SPONGIA.....	1-2 minim, 1 minim.....	75
SPONGIA.....	2 minims.....	1 15
STICTA.....	1 minim.....	75
THUJA.....	1 minim.....	75
THUJA.....	2 minims.....	1 15
URTICA URENS.....	1 minim.....	1 15
VERATRUM ALB.....	1-2 minim, 1 minim.....	75
VERATRUM ALB.....	2 minims.....	1 15
VERATRUM VIR.....	1-2 minim, 1 minim.....	75
VERATRUM VIR.....	2 minims.....	1 15
VIBURNUM OPULUS.....	1 minim.....	75
VIBURNUM PRUNIFOL.....	1 minim.....	75

TINCTURES.


	Per Pint.
TINCT. ACONITE.....	75
" ACONITE (Fleming's).....	90
" ARNICA.....	60
" BENZOIN COMP.....	80
" BRYONIA.....	70
" CANNABIS INDICA.....	1 50
" CANTHARIDES.....	75
" CAPSICUM.....	50
" CARDAMOM COMP.....	70
" CUBEBS.....	60
" CIMICIFUGA.....	70
" CINCHONA COMP.....	75
" DIGITALIS.....	75
" GENTIAN COMP.....	60
" GELSEMIUM.....	65
" GREEN SOAP.....	65
" GREEN SOAP COMP.....	85
" GINGER.....	95
" HYOSCYAMUS.....	50
" IGNATIA.....	85
" IODINE.....	1 00
" IODINE DECLORIZED.....	1 00
" IODINE COMP.....	1 25
" IODINE (Churchill's).....	1 50
" NUX VOMICA.....	75
" OPIUM, 10 per cent. of Opium.....	1 50
" OPIUM CAMPHORATED (Paregoric).....	50
" PHOSPHORUS.....	1 00
" SANGUINARIA.....	60
" STROPHANTHUS (Fraser).....	1 00

FRASER'S

Soluble Hypodermic Tablets.

The medicines in these tablets are combined with an unobjectionable base. They do not cause abscesses, and may be given internally as well as administered hypodermically.

These tablets are easily soluble in ten minims of warm water*; are hard enough to retain their form, and will not pulverize or crumble in the bottle. They also possess the advantage of accuracy of dose and perfect preservation of the drug.



Per Hundred.

ACONITINE NITRATE.....	1-50 gr.....	75
ACONITINE NITRATE.....	1-100 gr.....	60
ACONITINE NITRATE.....	1-150 gr.....	50
ACONITINE NITRATE.....	1-200 gr.....	40
APOMORPHINE MUR.....	1-10 gr.....	75
APOMORPHINE MUR.....	1-20 gr.....	55
ARSENITE OF POTASS	1-50 gr., 1-30 gr., 1-20 gr. .	30
ATROPINE SULPHATE.....	1-50 gr.....	35
ATROPINE SULPHATE .	1-200 gr., 1-150 gr., 1-100 gr.	30
BRUCIA NITRATE, (Merck's)...	1-100 gr., 1-50 gr.....	30
CAFFEINE CITRATE (Merck's)...	1-2 gr.....	40
CAFFEINE & SODIUM BENZ	1-2 gr.....	40
CAFFEINE & SODIUM BENZ	1 gr.....	60
COCAINE HYDROCHLORATE	1-20 gr.....	35
COCAINE HYDROCHLORATE	1-10 gr.....	55
COCAINE HYDROCHLORATE	1-8 gr.....	95
COCAINE HYDROCHLORATE	1-5 gr.....	95
COCAINE HYDROCHLORATE	1-4 gr.....	1 10
CODEINE PHOSPHATE.....	1-4 gr.....	1 10
CODEINE SULPHATE.....	1-4 gr.....	75
CORROSIVE SUBLIMATE.....	1-50 gr., 1-30 gr., 1-20 gr	30
DIGITALIA.....	1-100 gr.....	75
ERGOTIN.....	1-10 gr.....	30
HUBBARD'S BROMIDES No. 1	{ Morphine Brom. 1-4 gr. } { Hyoscin Hydro. 1-100 gr. }	1 25
HUBBARD'S BROMIDES No. 2	{ Hyoscin Hydrob. 1-100 gr. } { Morphine Brom. 1-8 gr. }	1 00


*It is unnecessary to use warm water for the Morphine, Atropine or Hyoscyamia Tablets; but better to use hot water for the others.

HYOSцин COMP.....	{ Hyoscin Hydrob. 1-200 gr. }	
(Dr. Cutter.)	{ Atropine Sulph. 1-200 gr. }	95
	{ Morphine Sulph. 1-4 gr. }	
HYOSцин HYDROBROM.....	1-200 gr.....	75
HYOSцин HYDROBROM.....	1-100 gr.....	95
HYOSциAMIA SULPHATE.....	1-200 gr.....	50
(Merck's White Cryst.)		
HYOSциAMIA SULPHATE.....	1-100 gr.....	75
(Merck's White Cryst.)		
HYOSциAMIA SULPHATE.....	1-50 gr.....	1 50
(Merck's White Cryst.)		
HYOSциAMIA SULPHATE.....	1-25 gr.....	3 00
(Merck's White Cryst.)		
MORPHINE SULPHATE.....	1-2 gr.....	75
MORPHINE SULPHATE.....	1-3 gr.....	55
MORPHINE SULPHATE.....	1-4 gr.....	45
MORPHINE SULPHATE.....	1-10 gr., 1-8 gr., 1-6 gr.....	35
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-2 gr. }	
	{ Atropine Sulph. 1-120 gr. }	75
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-3 gr. }	
	{ Atropine Sulph. 1-150 gr. }	55
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-4 gr. }	
	{ Atropine Sulph. 1-120 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-4 gr. }	
	{ Atropine Sulph. 1-150 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-4 gr. }	
	{ Atropine Sulph. 1-180 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-4 gr. }	
	{ Atropine Sulph. 1-200 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-6 gr. }	
	{ Atropine Sulph. 1-150 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-6 gr. }	
	{ Atropine Sulph. 1-180 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-8 gr. }	
	{ Atropine Sulph. 1-200 gr. }	40
MORPHINE & ATROPINE....	{ Morphine Sulph. 1-8 gr. }	
	{ Atropine Sulph. 1-150 gr. }	40
MORPHINE PHTHALATE*.....	1-8 gr.....	75
MORPHINE PHTHALATE.....	1-4 gr.....	1 00
MORPHINE PHTHALATE & ATROPINE.....	{ Morph. Phthalate. 1-4 gr. }	
	{ Atropine Sulph. 1-150 gr. }	1 10
MORPHINE PHTHALATE & ATROPINE.....	{ Morph. Phthalate. 1-6 gr. }	
	{ Atropine Sulph. 1-180 gr. }	1 00
MORPHINE PHTHALATE & ATROPINE.....	{ Morph. Phthalate. 1-8 gr. }	
	{ Atropine Sulph. 1-200 gr. }	90
MORPHINE & HYOSциAMIA.....	{ Morphine Sulph. 1-4 gr. }	
	{ Hyoscy. Sulph. 1-100 gr. }	1 10
NITRYGLYCERIN.....	1-200 gr. 1-100 gr. 1-50 gr.....	30
NITROGLYCERIN & STRYCH-NINE.....	{ Nitrogl. glycerin. 1-100 gr. }	
	{ Strych. Sulph. 1-50 gr. }	40
PILOCARPINE MUR.....	1-10 gr.....	2 00

*Phthalate of Morphine is more soluble than the sulphate, and has been specially recommended for hypodermic use, causing less nausea and general Morphine disturbance.

PILOCARPINE MUR.	1-20 gr.	1 00
QUININE BISULPHATE.....	1-2 gr.....	30
QUININE & UREA BI-MUR. ...	2 1-2 grs.	75
(For Hypodermic use.)		
STRYCHNINE NITRATE	{ 1-100 gr., 1-60 gr., 1-50 gr., 1-40 gr., 1-30 gr., 1-25 gr., 1-20 gr.....	30
STRYCHNINE SULPHATE....	{ 1-200 gr. 1-100 gr. 1-60 gr. } 1-50 gr. 1-25 gr.....	30

SYRINGE FOR HYPODERMIC TABLETS.



This syringe is made by THE FRASER TABLET TRITURATE MFG. CO., for convenience in dissolving and using their Hypodermic Tablets. It is provided with a moveable end or cap to protect the packing. For use the cap is removed and the tablet dropped into the syringe, together with ten minims of water and shaken for a few moments, until the tablet is dissolved.

Alligator Case.....	Net \$2 50
Aluminum "	" 2 75
Morocco "	" 2 50
Nickel "	" 2 50
Wood "	" 2 00

Please specify Fraser's Tablets

MISCELLANEOUS.

The attention of physicians is respectfully directed to the list of standard preparations from our own laboratory, which we present with confidence as containing in each case a first class specimen of the drug it represents faithfully and skillfully compounded, and of uniform strength.

PILLS.

BLAUD'S, Silver Coated	3 gr.....	per 1000	\$2 50
" " "	3 gr.....	per 100	35
" " "	5 gr.....	per 1000	3 00
" " "	5 gr.....	per 100	45
EXT. CANNABIS INDICA.....	1-4 gr.	per 1000	5 00
" " "	1-4 gr.....	per 100	75

Perfectly soluble in the stomach.

These pills are made from the Extract of Cannabis, which has given us such a reputation for quality and uniformity of the drug that we ship preparations of it all over the world. Every sample of the Extract is tested clinically before we make a preparation of it.

SPECIAL PREPARATIONS.

(Formerly manufactured by FRASER & Co., Incorporated.)

FRASER'S CHAMOMILLA COMP. (8 oz.)	Per Doz.	\$10.00
-----------------------------------	----------	---------

(The retail price of this preparation is \$1.25 per bottle, containing 64 adult doses.)

CHAMOMILLA COMP. is a diffusible nerve tonic, from the formula of Dr. Thos. A. McBride, and is a combination of the mother tinctures of Cinchona, Chamomilla, Ignatia and Phosphorus with Aromatics.

Samples of Chamomilla Comp. and literature referring to its use will be forwarded upon application.

FRASER'S GRANULES.

	Per Doz.	Per Gross.
PEPSIN AND SODA MINT GRANULES,	\$1.00	\$12.00
RHUBARB AND SODA MINT GRANULES,	1.00	12.00
SODA MINT GRANULES, - - -	85	10.00
CAMPBOR GLOBULES, - - -	1.00	12.00

THESE PREPARATIONS ARE PUT UP IN SCREW-CAP BOTTLES OF PATENTED DESIGN, and are packed in DOZENS for COUNTER DISPLAY, with a neat advertising card with each dozen. They are superior to any similar preparations ever offered.

Please specify **FRASER'S** in ordering or prescribing.

POWDER PAPERS.

POWDER PAPERS, No. 1, $2\frac{1}{2} \times 3\frac{1}{8}$	Per 1000.....	30
" " No. 2, $2\frac{3}{4} \times 3\frac{7}{8}$	" "	45
" " No. 3, $3\frac{5}{8} \times 5\frac{1}{8}$	" "	65

FRASER'S PIPETTES.



Fig. 2.


PLAIN, 4 inch.....	Each.....	Fig. 2	10
" 7 inch.....	"	"	10
" 12 inch.....	"	"	10
GRADUATED, 4 inch, tested. . .	20 minims.		10
" 7 " " . . .	30 "		25
" 12 " " . . .	40 "		25



Fig. 3.

GRADUATED, 4 inch, with Rubber suction, 20 min. Fig. 3.	20
" 7 " " " " 30 min. "	25
" 12 " " " " 40 min. "	30

DR. KEYS' ACID APPLICATOR.



Price.....	10
------------	----


URINE TESTS.

These tests are sold in two ounce glass stoppered bottles, and their reliability is guaranteed.

ACETIC ACID.....	25
CARBOL-FUCHSIN SOLUTION, For Stain	25
FEHLING'S TEST, Alkaline Solution.....	25
FEHLING'S TEST, Copper Solution	25
FRENCH CHALK, Prepared.....	25
INDIGO-CARMINE, Acidulated Solution.	25
INDIGO-CARMINE, Sodium Solution	25
MILLARD'S TEST.....	25
METHYLENE BLUE SOLUTION, For Stain.	25
SULPHURIC ACID, For Stain.....	25
TANRET'S TEST.....	25

Please specify FRASER'S in ordering or prescribing.



BOTTLES FOR DISPENSING TABLETS.

No. 50.
Full SizeNo. 25.
Full Size.No. 10.
Full Size.No. 50 A.
Full Size.No. 50 B.
Full Size.

THE NUMBERS OF THESE BOTTLES CORRESPOND TO THE NUMBER OF TABLETS EACH BOTTLE HOLDS. FOR INSTANCE, NO. 50 HOLDS ABOUT 50 TABLETS.

		Per Gross.
No. 50,	without corks	\$1 10
" 50,	with "	1 50
" 25,	without "	1 00
" 25,	with "	1 40
" 10,	without "	60
" 10,	with "	1 25
" 50A,	without "	1 10
" 50A,	with "	1 50
" 25A,	without "	1 00
" 25A,	with "	1 40
" 10A,	without "	60
" 10A,	with "	1 25
" 50B,	without "	1 10
" 50B,	with "	1 50
" 25B,	without "	1 00
" 25B,	with "	1 40
" 10B,	without "	60
" 10B,	with "	1 25

Please specify FRASER'S in ordering or prescribing.

PAPER DISPENSING VIALS.**No. 50,
Full Size.****No. 25,
Full Size.****No. 15,
Full Size.****No. 10,
Full Size.**

THE NUMBERS OF THESE VIALS CORRESPOND TO THE NUMBER OF TABLETS EACH VIAL HOLDS. THEY ARE PACKED IN PAPER, HALF GROSS RED AND HALF GROSS WHITE IN EACH PACKAGE.

No. 50.	Per Gross, \$1 00
" 25.	" " 75
" 15.	" " 70
" 10.	" " 60


STOCK CABINETS.

For Glass Stoppered Bottles of 1000 tablets.




No. 1.—Office Cabinet, to hold 25 Bottles..... \$2 00

Please specify **FRASER'S** in ordering or prescribing.



No. 2.—Revolving Cabinet, to hold 98 Bottles..... \$8 00



No. 3.—Wall Cabinet, to hold 108 Bottles..... 7 00
 No. 4. " " " " 192 " \$10 00

For.....
 Take..... Tablets

 Dr.

DIRECTION LABELS.

For Dispensing Bottles or Boxes
 printed like illustration and gummed.

Price, per 1000..... 50

Please specify **FRASER'S** in ordering or prescribing.

Vial Cases for Physicians.

No. 1.—Folding Pocket Emergency Case, $4\frac{1}{4}$ in. long by $3\frac{1}{2}$ in. wide when closed. Black Morocco and Seal Leather. 1 row of 12 tube vials, each holding 18 Tablet Triturates.

Price empty, - - - - - \$1 50
Filled with tablets as per following list, - - - - - 2 00

Aconite Tr., 1 minim.	Calomel, 2 grs.
Aloin and Belladonna Comp.	Digitalis Tr., 2 minims.
Ammonium Muriate Comp.	Dover's Tablets, 2 1-2 grs.
Belladonna Tr., 1 minim.	Morphine Sulphate, 1-4 gr.
Bronchitis.	Tartar Emetic, 1-50 gr.
Calomel, 1-10 gr.	Turpeth Mineral, 2 grs.

No. 2.—Folding Pocket Case with clasp. $7\frac{3}{4}$ in. long by $3\frac{1}{2}$ in. wide when closed. Black Seal Leather. One row 20 tube vials, each holding 18 Tablet Triturates.

Price empty, - - - - - \$2 50
Filled with any selection of medium-priced triturations, - - - - - 3 00

No. 3.—Folding Pocket Case with clasp. $7\frac{3}{4}$ in. long by $3\frac{1}{2}$ in. wide when closed. Black Seal Leather, with two rows of 20 tube vials, each holding 18 Tablet Triturates.

Price empty, - - - - - \$3 00
Filled with any selection of medium-priced triturations, - - - - - 3 75

No. 4.—Folding Pocket Case. $6\frac{1}{2}$ in. long by $3\frac{1}{2}$ in. wide when closed. Black Seal Leather, containing one row of 10 tube vials, each holding 40 Tablet Triturates.

Price empty, - - - - - \$2 50
Filled with any selection of medium priced tablets, - - - - - 3 00

No. 5.—Folding Pocket Case with clasp. $7\frac{3}{4}$ in. long by $3\frac{1}{2}$ in. wide when closed. Black Seal Leather, containing one row of 12 tube vials, each holding 40 Tablet Triturates.

Price empty, - - - - - \$2 50
Filled with any selection of medium-priced tablets, - - - - - 3 00



No. 6.—Folding Pocket Case with clasp. $7\frac{1}{2}$ in. long by $3\frac{1}{2}$ in. wide. Black Seal Leather, containing two rows of 12 tube vials, each vial holding 40 Tablet Triturates.

Price empty, - - - - - \$3 00
Filled with any selection of medium-priced tablets, - - - - - \$4 00



No. 7.—Hand Case. $9\frac{1}{2}$ in. long by $3\frac{1}{2}$ in. deep and $2\frac{1}{4}$ in. thick. Black or Russet Leather, with handle and lock, seamless patent round corners, containing 2 rows of 8 half-oz. bottles, each holding 100 Tablet Triturates.


Price empty, - - - - - \$5 00

Special Cases in any style Made To Order.




No. 8.—Buggy Case. $9\frac{1}{2}$ in. long by $6\frac{1}{2}$ in. deep and $2\frac{1}{4}$ in. thick. Black or Russet leather, with handle and lock, seamless patent round corners, containing four rows of 8 half-ounce bottles, each holding 100 Tablet Triturates.

Price empty, \$7 50



No. 9.—Dr. Hunter's Buggy Case. $10\frac{1}{2}$ in. long by $6\frac{1}{2}$ in. deep and $3\frac{1}{2}$ in. thick. Black or Russet leather, with handle and lock, patent round corners, 3 flaps inside, each flap containing two rows of 11 neck vials, making 66 vials, each holding 50 Tablet Triturates.


Price empty, \$12 00



No. 10.—Buggy Case. $10\frac{1}{2}$ in. long by 4 in. deep and 3 in. thick. Black Seal leather. 3 rows of 15 bottles and 1 row of 12 bottles and syringe case, with hypodermic syringe, 2 needles and 4 vials filled with hypodermic tablets. Case containing 57 tube vials, each holding about 50 Tablet Triturates.


Price with syringe complete and empty bottles, \$7 50

Special Cases in any style Made To Order.



No. 11.—Buggy Case, traveling bag pattern. 10 in. long, $6\frac{1}{2}$ in. deep, $5\frac{1}{2}$ in. thick. Black Seal Leather, with handle, lock and side clasps. Containing 7 rows of 8 half-ounce bottles and compartment $8 \times 2 \times 1$ in. for instruments, also pockets for dispensing boxes, etc.

Price empty, - - - - - \$12 50



No. 12.—Buggy Case. $9\frac{1}{2}$ in. long, $6\frac{1}{4}$ in. deep, 5 in. thick. Black leather with handle and double locks; seamless patent round corners, and contains 8 rows of 8 half-ounce screw top bottles, each capable of holding 100 tablet triturates.

Price empty, - - - - - \$13 50

Special Cases in any style Made to Order.



No. 13.—Tablet and Instrument Bag Size $13\frac{1}{2}$ in. long by $6\frac{1}{2}$ in. deep and 5 in. thick. Black Leather, contains 9 one ounce, 16 three ounce, and 20 two-ounce bottles on one side; the other side being an empty space entirely cut off from the medicine, for gynæcological or other instruments. This is a very useful bag for either city or country physicians.

Price empty, - - - - -

\$11 25



DR. STIMSON'S CASE.—Folding Leather Case with Clasps and Square Ends. Filled with $\frac{1}{2}$ or 1 oz. cork or screw cap bottles.

No. 1- 6 Bottles, Single Row,	\$1 50	No. 5-12 Bottles, Double Row,	\$3 50
No. 2- 8 " " "	2 00	No. 6-16 " " "	3 75
No. 3-10 " " "	2 50	No. 7-20 " " "	4 00
No. 4-12 " " "	3 00	No. 8-24 " " "	4 25

NOTE.—Any of our cases in which the bottles are held by clasps will be fitted with either leather loops or metal springs as may be desired. Unless otherwise specified we usually furnish leather loops.

Special Cases in any style Made to Order.

THE name "FRASER'S TABLETS" is recognized by medical practitioners everywhere as the synonym for tablet triturates, which were first developed, manufactured and brought to their present state of perfection by us, upon the suggestion of DR. ROBERT M. FULLER, (See *Medical Record*, March 9th, 1878 and March 25th, 1882.)

This method of dispensing medicines that are given in small quantities, has been attended, since its introduction by us, by the success due :

1st. To the care which we have taken in selecting and manufacturing the drugs and chemicals employed.

2d. To the thoroughness of the trituration and division of the remedy and consequent perfect accuracy of the dose.

3d. To the quickness and ease with which our tablets are dissolved or diffused in water and in the acids of the stomach.

4th. To their elegance of form and novelty of appearance.

CAUTION.

The manufacture of Tablet Triturates is apparently so simple that many have attempted to imitate our goods. We desire, therefore, to warn practitioners that dangers are possible in the use of tablets which may resemble ours, but are made without the painstaking care and skill, the high class drugs, the close adherence to the formula and above all the thorough trituration, which lead to the uniformity of product that has been a recognized characteristic of our goods from the first. In prescribing the small doses represented by many of the preparations in tablet triturate form, these characteristics are indispensable to the success of the remedies.

